Jim de Groot

College of Engineering and Computer Science The Australian National University Canberra, ACT, Australia jim.degroot@anu.edu.au

Abstract

We prove Goldblatt-Thomason theorems for frames and models of a wide variety of modal intuitionistic logics, including ones studied by Wolter and Zakharyaschev, Goldblatt, Fischer Servi, and Plotkin and Sterling. We use the framework of dialgebraic logic to describe most of these logics and derive results in a uniform way.

1 Introduction

A prominent question in the study (modal) logics and their frame semantics is what classes of frames can be defined as the class of frames satisfying some set of formulae. Such a class is usually called *axiomatic* or *modally definable*. A milestone result partially answering this question in the realm of classical normal modal logic is a theorem by Goldblatt and Thomason [14] and dates back to 1974. It states that an elementary class of Kripke frames is axiomatic if and only if it reflects ultrafilter extensions and is closed under bounded morphic images, generated subframes and disjoint unions. The proof given in *op. cit.* relies on Birkhoff's variety theorem and makes use of the algebraic semantics of the logic. A model-theoretic proof was provided almost twenty years later by Van Benthem [33].

Analogous results for classes of *models*, rather than frames, are also of interest. In case of classical normal modal logic these were investigated in the nineties by Barwise and Moss [1], and Venema [34]. Such theorems, both for frames and for models, are nowadays often referred to as *Goldblatt-Thomason theorems*.

A similar result for (non-modal) intuitionistic logic was proven by Rodenburg [28] (see also [13]), where the interpreting structures are *intuitionistic* Kripke frames and models. This, of course, requires analogs of the notions of bounded morphic images, generated subframes, disjoint unions and ultrafilter extensions. While the first three carry over straightforwardly from the setting of classical normal modal logic, ultrafilters have to be replaced by *prime filters*.

In recent years, Goldblatt-Thomason theorems for many other logics have been proven as well, including ones for positive normal modal logic [5], graded modal logic [30], modal extensions of Lukasiewicz finitely-valued logics [32], LE-logics [8], and several versions of modal logic with a universal modality [31].

A more general approach towards Goldblatt-Thomason theorems, using the framework of coalgebraic logic, was taken by Kurz and Rosický [22]. Coalgebraic logic is a framework in which modal logics are developed parametric in the signature of the language and an endofunctor $\mathcal{T} : C \to C$ on some base category C (see e.g. [25, 21] for an introduction). In [22], the authors focus on the case where \mathcal{T} is an endofunctor on Set, and prove Goldblatt-Thomason theorems for a wide variety of such \mathcal{T} . In particular, the theorem holds when \mathcal{T} is a generalised Kripke polynomial functor, a notion that originates from [29]. These are functors obtained from

the identity functor, constant functors, the powerset functor and the neighbourhood functor, by combining them using products, coproducts and composition. Each generalised Kripke polynomial functor comes with a modal logic for which a Goldblatt-Thomason theorem is given.

In the present paper we prove Goldblatt-Thomason theorems for a number of modal *intuitionistic* logics. Amongst these are logics by Wolter and Zakharyaschev [35, 36, 37], modal logic with a geometric modality [12, Section 6] and the modal intuitionistic logic studied by Fischer Servi [10, 11] and Plotkin and Sterling [26]. For each of these logics we obtain:

> A class K of frames closed under prime filter extensions is axiomatic if and only if it reflects prime filter extensions and is closed under disjoint unions, regular subframes and p-morphic images.

and

A class M of models is axiomatic if and only if it is invariant under prime filter extensions, and closed under disjoint unions, p-morphic domains and p-morphic images.

Evidently, the definitions of frame, model, regular subframe, p-morphism, disjoint union and prime filter extension depend on the logic and semantics under consideration.

The main technical tool we use to describe most of these logics is the framework of *dialgebraic* logic. This is a framework where modal logics are developed parametric in the signautre of the language and a functor $\mathcal{T} : \mathbb{C}' \to \mathbb{C}$, where \mathbb{C}' is some subcategory of \mathbb{C} . It slightly generalises coalgebraic logic and was recently introduced in [17]. While coalgebraic logics are too restrictive to describe modal intuitionistic logics (see e.g. [23, Remark 8] or [17, Section 2] for a more detailed discussion), the additional flexibility of dialgebraic logic does allow one to model a number of them. In particular, modal intuitionistic logics studied in [12, 35, 37] fit this framework. In Section 2 we recall the required prerequisites of dialgebraic logic and illustrate these with examples, so that no prior knowledge of dialgebraic logic is required.

Our strategy for obtaining Goldblatt-Thomason theorems for such *dialgebraic intuitionistic logics* closely resembles the approach taken by Goldblatt and Thomason in [14]. The framework of dialgebraic logic comes with its built-in notion of algebraic semantics, to which we can apply Birkhoff's variety theorem. If a dialgebraic intuitionistic logic satisfies a certain *canonicity* condition, we are guaranteed a notion of prime filter extension. We then use these prime filter extensions to transfer the variety theorem on the algebraic semantics to a Goldblatt-Thomason theorem for the frame semantics.

To illustrate that there exists a myriad of canonical dialgebraic intuitionistic logics, we define the class of *intuitionistic polynomial functors*, each of which comes equipped with a logic that is canonical. These functors are built from certain variations of the identity functor, constant functor, powerset functor and neighbourhood functor, that can be then combined using products, coproductss, and a variation of coproducts. As a consequence of our main results, we obtain Goldblatt-Thomason theorems for the frames and models of each of these dialgebraic intuitionistic logics.

The modal intuitionistic logic studied by Fischer Servi [10, 11] and Plotkin and Sterling [26] is *not* a dialgebraic intuitionistic logic. But there is no need for sadness over this: in Section 7 we show how to obtain a Goldblatt-Thomason theorem for this logic using our general results.

Structure of the paper

In Section 2 we recall the framework of a dialgebraic logic specified to the setting of intuitionistic logic. We give the definition of frame semantics, algebraic semantics, canonicity and prime filter extensions, and illustrate each of these by concrete examples from [35, 12]. Subsequently, in Section 3, we give four more examples of modal intuitionistic logics in a dialgebraic format. The functors corresponding to these logics will serve as the "atoms" of our notion of intuitionistic polynomial functor.

In Section 4 we give three ways of combining functors \mathcal{T} : Krip \rightarrow Pos and their logics: using products, coproducts, and a variation of coproducts. Together with the functors defined in Sections 2 and 3 these operations give rise to the class of *intuitionistic polynomial functors*. We also prove that each of these functors comes with a canonical modal intuitionistic logic.

Section 5 is devoted to invariance results: we prove that validity of formulae is preserved under certain notions of morphisms, bisimulations, disjoint unions, and prime filter extensions. This is, of course, one implication of the Goldblatt-Thomason theorem. Thereafter, in Section 6, we complete the proof of the Goldblatt-Thomason theorem for canonical dialgebraic intuitionistic logics.

Finally, in Section 7 we prove the Goldblatt-Thomason theorem for the modal intuitionistic logic studied by Fischer Servi [10, 11] and Plotkin and Sterling [26] using the general result from Section 6.

2 A framework for modal intuitionistic logics

We recall the framework of dialgebraic logic, specified to the case of modal intuitionistic logics. This framework was recently introduced in [17] and the contents of this Section have appeared in [17] in a condensed form with the following exception: Our treatment of proposition letters is different. While it is noted in *op. cit.* that proposition letters can be introduced via predicate liftings, we "manually" add them to our logic. This leads to a notion of model on top of a dialgebra. The reason for this change is twofold: first, it makes it easier to relate the abstract framework to concrete examples; second, from an algebraic point of view proposition letters behave like variables, and their presence allows us to prove a Goldblatt-Thomason theorem in a relatively straightforward manner.

Throughout this section, we connect the abstract definitions to two concrete examples: intuitionistic logic with a normal modal box from [37] and intuitionistic logic with a monotone modal operator from [12, Section 6]. Before we get started, we fix notation for frequently-used categories and functors in Subsection 2.1

2.1 Some categories and functors

We denote posets by X, X', Y, \ldots and suppress the order. If the order is relevant then we denote it by $\leq_X, \leq_{X'}, \leq_Y$. A monotone function from X to X' is a map $f: X \to X'$ that preserves the order, i.e., $x \leq_X y$ implies $f(x) \leq_{X'} f(y)$. Let **Pos** be the category of posets and monotone functions. An *embedding* is a function $f: X \to X'$ that preserves and reflects the poset order, that is, $x \leq_X y$ if and only if $f(x) \leq_{X'} f(y)$.

A monotone function $f: X \to X'$ is called *bounded* if for all $x \in X$ and $y' \in X'$, $f(x) \leq_{X'} y'$ implies the existence of some $y \in X$ such that $x \leq_X y$ and f(y) = y'. We write Krip for the category of posets and bounded morphisms, and $i: \text{Krip} \to \text{Pos}$ for the inclusion functor.

Furthermore, we let DL be the category of distributive lattices and homomorphisms, and HA the category of Heyting algebras and Heyting algebra homomorphisms. Let $j : HA \rightarrow DL$ be the obvious inclusion functor.

Recall that the collection of upwards-closed subsets of a poset X forms a distributive lattice. This assignment extends to a contravariant functor $up : \text{Pos} \to \text{DL}$, with $upf = f^{-1}$ for morphisms $f \in \text{Pos}$. This, in turn, restricts to a functor $\text{Krip} \to \text{HA}$ that we denote by up'. In the converse direction, we have a contravariant functor $pf : \text{DL} \to \text{Pos}$ that sends a distributive lattice A to the set of prime filters of A ordered by inclusion, and a homomorphism $h : A \to A'$ to $h^{-1} : pfA' \to pfA$. (In other words, pf takes the Priestley [27] and then forgets about the topology.) This functors restricts to $pf' : \text{HA} \to \text{Krip}$. The functors we have just introduced yield the following (commuting) diagrams:

$$\begin{array}{ccc} \operatorname{Pos} & \stackrel{up}{\longrightarrow} & \operatorname{DL} & \operatorname{Pos} & \stackrel{pf}{\longleftarrow} & \operatorname{DL} \\ i & \uparrow & \uparrow^{j} & i & \uparrow^{j} \\ \operatorname{Krip} & \stackrel{up'}{\longrightarrow} & \operatorname{HA} & \operatorname{Krip} & \stackrel{pf'}{\longleftarrow} & \operatorname{HA} \end{array}$$

The functors up and pf constitute a dual adjunction between Pos and DL. The units

$$\eta : \mathrm{id}_{\mathsf{Pos}} \to pf \cdot up, \qquad \theta : \mathrm{id}_{\mathsf{DL}} \to up \cdot pf,$$

of this dual adjunction are defined by $\eta_X(x) = \{a \in upX \mid x \in a\}$ and $\theta_A(a) = \tilde{a} = \{p \in pfA \mid a \in p\}$. If A is a Heyting algebra, then $\theta_A : A \to up(pfA)$ is a Heyting algebra homomorphism (this follows from Esakia duality, see e.g. [9]). We write $\theta' : id_{\mathsf{HA}} \to up' \cdot pf'$ for the restriction of θ to HA. (In the language of [17, Definition 5.2], the existence of this θ' means that our setup is well-structured.)

2.2 Dialgebras

We recall the definition of a dialgebra, and give two examples of interpreting structures for modal intuitionistic logics from the literature, recast as a category of dialgebras.

2.1 Definition. Let $\mathcal{F}, \mathcal{G} : \mathsf{C} \to \mathsf{D}$ be a parellel pair of functors from a category C to a category D . An $(\mathcal{F}, \mathcal{G})$ -dialgebra is a pair (X, γ) of an object X in C and a morphism $\gamma : \mathcal{F}X \to \mathcal{G}X$ in D . An $(\mathcal{F}, \mathcal{G})$ -dialgebra morphism $f : (X, \gamma) \to (X', \gamma')$ is a morphism $f : X \to X'$ in C satisfying $\mathcal{G}f \circ \gamma = \gamma' \circ \mathcal{F}f$. In diagrams,



We denote the category of $(\mathcal{F}, \mathcal{G})$ -dialgebras and $(\mathcal{F}, \mathcal{G})$ -dialgebra morphisms by $\mathsf{Dialg}(\mathcal{F}, \mathcal{G})$.

In this paper we focus on two types of dialgebras. First, we are interested in dialgebras for functors $\mathcal{T} : \mathsf{Krip} \to \mathsf{Pos}$. In particular, (i, \mathcal{T}) -dialgebras will play the rôle of frame semantics of modal intuitionistic logics (recall that $i : \mathsf{Krip} \to \mathsf{Pos}$ is the inclusion functor). Therefore we will sometimes refer to (i, \mathcal{T}) -dialgebras as \mathcal{T} -dialgebras or \mathcal{T} -frames. Besides, in analogy with modal logic, we sometimes call a (i, \mathcal{T}) -dialgebra morphism a *p*-morphism.

Second, we will see dialgebras for functors $\mathcal{L}, j : HA \to DL$, where j is the inclusion of categories. Such dialgebras help us to abstractly represent the algebraic semantics of a wide variety of modal intuitionistic logics.

We now list two examples of (categories of) dialgebras. These will serve as (the state spaces of) the running examples of this section.

2.2 Example. A \Box -frame is a tuple (X, R) consisting of a poset X and a relation R that satisfies

$$(\leq_X \circ R \circ \leq_X) = R.$$

A morphism between \Box -frames is a function between the underlying sets that is bounded with respect to both the poset order as well as R. We write $WZ\Box$ for the category of \Box -frames and \Box -frame morphisms. These frames are used in e.g. [37] to interpret intuitionistic logic with a box-modality and also make an appearance in [38, Definition 1.1.5] and [19, Definition 1].

The category WZD can be described as a category of dialgebras for the functors i, \mathcal{P}_{up} : Krip \rightarrow Pos, i.e., WZD \cong Dialg (i, \mathcal{P}_{up}) . Here \mathcal{P}_{up} sends a poset X to its collection of up-closed subsets ordered by reverse inclusion, and a bounded morphism $f: X \rightarrow X'$ to the direct image function $\mathcal{P}_{up}f = f[-]: \mathcal{P}_{up}X \rightarrow \mathcal{P}_{up}X'$. For details we refer to [17, Section 2].

2.3 Example. An *intuitionistic monotone frame*, or *monotone frame* for short, is a pair (X, N) where X is a poset and N is a function that selects a collection of neighbourhoods $N(x) \subseteq upX$ for every $x \in X$ such that:

- If $a \in N(x)$ and $a \subseteq b \in upX$ then $b \in N(x)$;
- If $x \leq y$ then $N(x) \subseteq N(y)$

Such frames are called *neighbourhood spaces* by Goldblatt in [12, Section 6.4.1], who uses them as interpreting structures for intuitionistic logic with a monotone modality. Although not defined in *op. cit.*, the obvious notion of a morphism between monotone frames (X, N) and (X', N') is a bounded morphism $f: X \to X'$ such that for all $x \in X$ and $a' \in up'X'$:

$$f^{-1}(a') \in N(x)$$
 iff $a' \in N'(f(x))$.

With this notion of morphism, monotone frames form a category that we denote by Mon.

As observed in [17, Section 8], the category Mon is isomorphic to a category of dialgebras. Define $\mathcal{H} : \mathsf{Krip} \to \mathsf{Pos}$ on objects by sending a poset X to

$$\mathcal{H}X = \{W \subseteq upX \mid \text{If } a \in W \text{ and } a \subseteq b \in upX \text{ then } b \in W\}$$

ordered by inclusion. If $f: X \to X'$ is a morphism in Krip, define $\mathcal{H}f$ by $\mathcal{H}f(W) = \{a' \in upX' \mid f^{-1}(a') \in W\}$. Then

Mon
$$\cong$$
 Dialg (i, \mathcal{H}) .

2.3 Interpreting modalities

It is well known that intuitionistic formulae can be interpreted in posets with a valuation. We now present a method of defining modal operators that can be interpreted in (i, \mathcal{T}) -dialgebras, where $i : \text{Krip} \to \text{Pos}$ is the inclusion functor and $\mathcal{T} : \text{Krip} \to \text{Pos}$ is any functor. This is a straightforward adaptation of the notion of a predicate lifting from the setting of coalgebraic logic for set-coalgebras [25], and was presented in [17, Section 5].

We write $\mathcal{U}p : \mathsf{Pos} \to \mathsf{Set}$ for the contravariant functor that sends a poset X to the set of upwards closed subsets of X, and a monotone function $f : X \to X'$ to the inverse images $\mathcal{U}pf = f^{-1} : \mathcal{U}pX' \to \mathcal{U}pX$. In other words, $\mathcal{U}p$ is the composition of the functors $up : \mathsf{Pos} \to \mathsf{DL}$ and the forgetful functor $\mathcal{U} : \mathsf{DL} \to \mathsf{Set}$.

2.4 Definition. An *n*-ary predicate lifting for (i, \mathcal{T}) -dialgebras is a natural transformation

$$\lambda: \mathcal{U}p^n \cdot i \to \mathcal{U}p \cdot \mathcal{T},$$

where $\mathcal{U}p^n X$ is the *n*-fold product of $\mathcal{U}pX$ in Set.

Throughout this paper, we fix an infinite set Prop of proposition letters. A collection Λ of predicate liftings for (i, \mathcal{T}) -dialgebras gives rise to the language $\mathbb{L}(\Lambda)$ given by the grammar

$$\varphi ::= p \mid \top \mid \bot \mid \varphi \land \varphi \mid \varphi \lor \varphi \mid \varphi \to \varphi \mid \heartsuit^{\lambda}(\varphi_1, \dots, \varphi_n)$$

where p ranges over Prop and $\lambda \in \Lambda$ is an n-ary predicate lifting. This language can be interpreted in \mathcal{T} -models:

2.5 Definition. A valuation for a \mathcal{T} -frame (X, γ) is an assignment $V : \operatorname{Prop} \to \mathcal{U}pX$ of the proposition letters. A \mathcal{T} -model consists of a \mathcal{T} -frame together with a valuation. The interpretation of $\mathbb{L}(\Lambda)$ -formulae in a state x of a \mathcal{T} -model $\mathfrak{M} = (X, \gamma, V)$ is defined recursively via

$$\begin{split} \mathfrak{M}, x \Vdash p & \text{iff} \quad x \in V(p) \\ \mathfrak{M}, x \Vdash \top & \text{always} \\ \mathfrak{M}, x \Vdash \bot & \text{never} \\ \mathfrak{M}, x \Vdash \varphi \land \psi & \text{iff} \quad \mathfrak{M}, x \Vdash \varphi \text{ and } \mathfrak{M}, x \Vdash \psi \\ \mathfrak{M}, x \Vdash \varphi \lor \psi & \text{iff} \quad \mathfrak{M}, x \Vdash \varphi \text{ or } \mathfrak{M}, x \Vdash \psi \\ \mathfrak{M}, x \Vdash \varphi \lor \psi & \text{iff} \quad x \leq y \text{ and } \mathfrak{M}, y \Vdash \psi \\ \mathfrak{M}, x \Vdash \varphi \to \psi & \text{iff} \quad x \leq y \text{ and } \mathfrak{M}, y \Vdash \varphi \text{ imply } \mathfrak{M}, y \Vdash \psi \\ \mathfrak{M}, x \Vdash \varphi^{\lambda}(\varphi_{1}, \dots, \varphi_{n}) & \text{iff} \quad \gamma(x) \in \lambda_{X}(\llbracket \varphi_{1} \rrbracket^{\mathfrak{M}}, \dots, \llbracket \varphi_{n} \rrbracket^{\mathfrak{M}}) \end{split}$$

Here $\llbracket \varphi \rrbracket^{\mathfrak{M}} = \{x \in X \mid x \Vdash \varphi\}$ denotes the *truth set* of φ . If no confusion is likely we sometimes write $\llbracket \varphi \rrbracket$ instead of $\llbracket \varphi \rrbracket^{\mathfrak{M}}$.

A model $\mathfrak{M} = (X, \gamma, V)$ is said to *satisfy* a formula φ if $\llbracket \varphi \rrbracket^{\mathfrak{M}} = X$, notation: $\mathfrak{M} \Vdash \varphi$. We say that \mathfrak{M} satisfies a set Φ of $\mathbb{L}(\Lambda)$ -formulae, notation: $\mathfrak{M} \Vdash \Phi$, if it satisfies every $\varphi \in \Phi$. A frame $\mathfrak{X} = (X, \gamma)$ is said to satisfy φ if $(X, \gamma, V) \Vdash \varphi$ for all valuations V (written as $\mathfrak{X} \Vdash \varphi$) and we similarly define $\mathfrak{X} \Vdash \Phi$.

The collection of \mathcal{T} -models forms a category with the following notion of morphism.

2.6 Definition. A morphism between \mathcal{T} -models (X, γ, V) and (X', γ', V') is an (i, \mathcal{T}) -dialgebra morphism $f: (X, \gamma) \to (X', \gamma')$ that satisfies

$$V' = V \circ \mathcal{U}pf.$$

We write $\mathsf{Mod}(\mathcal{T})$ for the category of \mathcal{T} -models and their morphisms.

2.7 Example. In [37] the \Box -frames from Example 2.2 are used to interpret intuitionistic logic with a \Box modality. As in classical modal logic, the interpretation of \Box is given by $x \Vdash \Box \varphi$

Jim de Groot

iff xRy implies $y \Vdash \varphi$. From the point of view of predicate liftings, this corresponds to λ^{\Box} : $\mathcal{U}p \cdot i \to \mathcal{U}p \cdot \mathcal{P}_{up}$ defined by

$$\lambda_X^{\square}(a) = \{ b \in \mathcal{P}_{up}X \mid b \subseteq a \}$$

We then have

 $x \Vdash \Box \varphi \quad \text{iff} \quad \gamma(x) \subseteq \llbracket \varphi \rrbracket$

and since $\gamma(x)$ denotes the set of successors of a state x, this definition corresponds to the intended one.

2.8 Example. The language \mathbb{L}_{Δ} is the extension of the intuitionistic grammar with a monotone modal operator Δ , called a *geometric modality* in [12]. Formulae in this language can be interpreted in a model (X, N, V) based on monotone frame (X, N). The intuitionistic connectives are interpreted in the underlying poset X as usual, and truth of Δ is defined by

$$x \Vdash \bigtriangleup \varphi$$
 iff $\llbracket \varphi \rrbracket \in N(x)$

We can obtain \triangle from the predicate lifting λ^{\triangle} defined on components by

$$\lambda_X^{\Delta}: \mathcal{U}p(iX) \to \mathcal{U}p(\mathcal{H}X): a \mapsto \{W \in \mathcal{H}X \mid a \in W\}$$

Identifying \triangle in the grammer of \mathbb{L}_{\triangle} with $\heartsuit^{\lambda^{\triangle}}$ in $\mathbb{L}(\{\lambda^{\triangle}\})$ then yields an obvious correspondence between \mathbb{L}_{\triangle} and $\mathbb{L}(\{\lambda^{\triangle}\})$, and in a monotone model (X, N, V) (viewed as an \mathcal{H} -model) we have

$$x \Vdash \heartsuit^{\lambda^{\Delta}} \varphi$$
 iff $N(x) \in \lambda^{\Delta}(\llbracket \varphi \rrbracket)$ iff $\llbracket \varphi \rrbracket \in N(x)$ iff $x \Vdash \bigtriangleup \varphi$.

2.4 Algebraic semantics

If the axioms for the extra modal operators have rank 1 and do not involve implication, then the algebraic semantics of the logic can be described via dialgebras. Let us make this more precise.

For a set S, we write $\mathcal{T}erS$ for the term algebra over S for the signature $\{\top, \bot, \land, \lor\}$. If Λ is a collection of predicate liftings and A is a set then we define

$$\Lambda(A) = \{ \heartsuit^{\lambda}(a_1, \dots, a_n) \mid \lambda \in \Lambda \text{ n-ary, } a_i \in A \}.$$

We underline $\underline{\heartsuit}^{\lambda}$ to indicate that they are formal objects rather than formulae in $\mathbb{L}(\Lambda)$. We can now define the notion of a Λ -axiom. Intuitively, these are axioms of modal depth 1 that do not involve intuitionistic implication.

2.9 Definition. A Λ -axiom is a pair (φ, ψ) of elements in $\mathcal{T}er(\Lambda(\mathcal{T}erS))$, where S is some set of variables.

From a logical point of view, a Λ -axiom gives rise to an axiom for the language $\mathbb{L}(\Lambda)$ by replacing the variables with proposition letters. This then yields a calculus $\mathbf{IPC}(\Lambda, Ax)$ for the language $\mathbb{L}(\Lambda)$, which extends \mathbf{IPC} with the axioms arising from the Λ -axioms in Ax, and with the congruence rule

$$\frac{p_1 \leftrightarrow q_1 \cdots p_n \leftrightarrow q_n}{\heartsuit^{\lambda}(p_1, \dots, p_n) \leftrightarrow \heartsuit^{\lambda}(q_1, \dots, q_n)}$$

The algebraic semantics of this logic are given as follows:

2.10 Definition. Let Λ be a set of predicate liftings for (i, \mathcal{T}) -dialgebras and Ax a collection of one-step axioms for Λ . Then we define $\mathcal{L} : \mathsf{HA} \to \mathsf{DL}$ on objects by sending a Heyting algebra A to the free distributive lattice generated by $\Lambda(A)$ modulo the relations from Ax, where the variables are substituted for elements in A. For a morphism $h : A \to A'$ in HA define $\mathcal{L}h$ on generators by

$$\mathcal{L}h(\underline{\heartsuit}^{\lambda}(a_1,\ldots,a_n))=\underline{\heartsuit}^{\lambda}(h(a_1),\ldots,h(a_n)).$$

The collection of predicate liftings now give rise to a transformation $\rho_{(\Lambda, Ax)} : \mathcal{L} \cdot up' \to up \cdot \mathcal{T}$ via

$$\rho_{(\Lambda,\mathrm{Ax}),X}([\heartsuit^{\lambda}(a_1,\ldots,a_n)]) = \lambda_X(a_1,\ldots,a_n),$$

where $[\cdot]$ denotes the equivalence class of a in $\mathcal{L}A$. We call Ax sound if $\rho_{(\Lambda, Ax)}$ is well defined. In this case, naturality of $\rho_{(\Lambda, Ax)}$ is an immediate consequence of naturality of the predicate liftings.

Abstractly, one can define a logic for (i, \mathcal{T}) -dialgebras as a pair (\mathcal{L}, ρ) where \mathcal{L} is a functor $\mathsf{HA} \to \mathsf{DL}$ and ρ a natural transformation $\mathcal{L} \cdot up' \to up \cdot \mathcal{T}$, see [17, Definition 5.3]. If such a pair arises via the procedure described above we say that (\mathcal{L}, ρ) is given by predicate liftings and axioms. In the current paper we will exclusively be interested in logics given by predicate liftings and axioms, because this implies that the category $\mathsf{Dialg}(\mathcal{L}, j)$ is a variety of algebras. The natural transformation $\rho_{(\Lambda, \mathrm{Ax})}$ proves useful when defining a property that we call canonicity in the next subsection.

The formulae from $\mathbb{L}(\Lambda)$ can be interpreted in (\mathcal{L}, j) -dialgebras with an assignment of the proposition letters.

2.11 Definition. Let (\mathcal{L}, ρ) be a logic for a functor \mathcal{T} given by a pair (Λ, Ax) of predicate liftings and axioms. An *algebraic model* is a tuple (A, α, ζ) where (A, α) is an (\mathcal{L}, j) -dialgebra and ζ : Prop $\to A$ is an assignment of the proposition letters. The function ζ extends to an interpretation of all $\mathcal{L}(\Lambda)$ -formulae via

$$\begin{split} \widetilde{\zeta}(p) &= \zeta(p) \\ \widetilde{\zeta}(\top) &= \top_A \\ \widetilde{\zeta}(\bot) &= \bot_A \\ \widetilde{\zeta}(\varphi \star \psi) &= \widetilde{\zeta}(\varphi) \star_A \widetilde{\zeta}(\psi) \\ \widetilde{\zeta}(\heartsuit^{\lambda}(\varphi_1, \dots, \varphi_n)) &= \alpha(\underline{\heartsuit}^{\lambda}(\widetilde{\zeta}(\varphi_1), \dots, \widetilde{\zeta}(\varphi_n))) \end{split}$$

Here \star_A indicates a connective viewed as an operation in the Heyting algebra A. If (A, α, ζ) is an algebraic model, then we write $(A, \alpha, \zeta) \models \varphi$ if $\tilde{\zeta}(\varphi) = \top_A$, and say that (A, α, ζ) satisfies φ . For $\Phi \subseteq \mathbb{L}(\Lambda)$ and $(A, \alpha) \in \mathsf{Dialg}(\mathcal{L}, j)$ define $(A, \alpha) \models \varphi$, $(A, \alpha, \zeta) \models \Phi$ and $(A, \alpha) \models \Phi$ in the standard way.

A morphism between algebraic models is a dialgebra morphism that preserves the assignment. We write $\mathsf{AlgMod}(\mathcal{L})$ for the category of algebraic models and their morphisms.

Every \mathcal{T} -frame $\mathfrak{X} = (X, \gamma)$ gives rise to an (\mathcal{L}, j) -dialgebra $\mathfrak{X}^+ = (up'X, \gamma^+)$, called the *complex algebra* of \mathfrak{X} . Here γ^+ is defined by the composition

$$\mathcal{L} \cdot up'X \xrightarrow{\rho_X} up \cdot \mathcal{T}X \xrightarrow{up\gamma} up \cdot iX = j \cdot up'X.$$

This assignment can be extended to a functor $(\cdot)^+$: $\text{Dialg}(i, \mathcal{T}) \to \text{Dialg}(\mathcal{L}, j)$ by setting $f^+ = up'f$ $(=f^{-1})$ for an (i, \mathcal{T}) -dialgebra morphism f. Valuations for a \mathcal{T} -frame (X, γ) correspond

bijectively with assignments for \mathfrak{X}^+ , and this yields a functor

$$(\cdot)^+ : \mathsf{Mod}(\mathcal{T}) \to \mathsf{AlgMod}(\mathcal{L}) : (X, \gamma, V) \mapsto (up'X, \gamma^+, V)$$

given on morphisms by $f^+ = up'f$ again.

By definition of the interpretation of formulae in (algebraic) models we have $\mathfrak{M} \Vdash \varphi$ if and only if $\mathfrak{M}^+ \models \varphi$ for all \mathcal{T} -models \mathfrak{M} . Moreover, since valuations for \mathfrak{X} correspond bijectively to assignments for \mathfrak{X}^+ we have $\mathfrak{X} \Vdash \varphi$ if and only if $\mathfrak{X}^+ \models \varphi$ for all \mathcal{T} -frames \mathfrak{X} .

2.12 Example. Consider the functor \mathcal{P}_{up} : $\mathsf{Krip} \to \mathsf{Pos}$ and the predicate lifting λ^{\Box} given by $\lambda^{\Box}_X(a) = \{b \in \mathcal{P}_{up}X \mid b \subseteq a\}$. Then the axioms

$$\underline{\Box}a \wedge \underline{\Box}b = \underline{\Box}(a \wedge b), \qquad \underline{\Box}\top = \top$$

are easily seen to be sound. The algebraic semantics that arise from λ^{\Box} and these axioms are given by (\mathcal{L}_{\Box}, j) -dialgebras, where \mathcal{L}_{\Box} sends a Heyting algebra A to the distributive lattice generated by $\{\underline{\Box}a \mid a \in A\}$ modulo the axioms. The corresponding natural transformation $\rho^{\Box}: \mathcal{L}_{\Box} \cdot up' \to up \cdot \mathcal{P}_{up}$ is defined on generators by $\rho_X^{\Box}(\underline{\Box}a) = \{b \in \mathcal{P}_{up}X \mid b \subseteq a\}.$

Alternatively, we can view (\mathcal{L}_{\Box}, j) -dialgebras as pairs (A, \Box) consisting of a Heyting algebras A and a finite meet-preserving map $\Box : A \to A$. A (\mathcal{L}_{\Box}, j) -dialgebra (A, α) gives rise to such a pair by setting $\Box a = \alpha(\underline{\Box}a)$. Conversely, given (A, \Box) we can define a dialgebra structure map α for A via $\alpha(\underline{\Box}a) = \Box a$.

2.13 Example. The monotone operator \triangle interpreted in monotone models satisfies the monotonicity axiom

$$\underline{\Delta}(a \wedge b) \leq \underline{\Delta}a.$$

In this case the algebraic semantics are given by $(\mathcal{L}_{\Delta}, j)$ -dialgebras, where \mathcal{L}_{Δ} sends a Heyting algebra A to the free distributive lattice generated by $\{\underline{\Delta}a \mid a \in A\}$ modulo monotonicity. The induced natural transformation $\rho : \mathcal{L}_{\diamond} \cdot up' \to up \cdot \mathcal{H}$ is given on generators by $\rho_X(\underline{\Delta}a) = \{W \in \mathcal{H}X \mid a \in W\}$.

2.5 Canonical logics and prime filter extensions

Our proof of the Goldblatt-Thomason theorem relies on Birkhoff's variety theorem and the connection between frame semantics and algebraic semantics of a logic. As we have seen above, in the general setup of dialgebraic modal logic we always have a functor $\text{Dialg}(i, \mathcal{T}) \rightarrow \text{Dialg}(\mathcal{L}, j)$. However, a converse is not naturally present. A sufficient condition is for the existence of a functor $\text{Dialg}(\mathcal{L}, j) \rightarrow \text{Dialg}(i, \mathcal{T})$ is the existence of a section of the adjoint buddy ρ^{\flat} of ρ , which is defined by

$$\rho^{\flat}: \mathcal{T} \cdot pf' \xrightarrow{\eta_{\mathcal{T}:pf'}} pf \cdot up \cdot \mathcal{T} \cdot pf' \xrightarrow{pf \rho_{pf'}} pf \cdot \mathcal{L} \cdot up' \cdot pf' \xrightarrow{pf \cdot \mathcal{L}\theta'} pf \cdot \mathcal{L}.$$

Recall that η is a unit of the dual adjunction between Pos and DL, and θ' is the restriction of the other unit $\theta : id_{\mathsf{DL}} \to up \cdot pf$ to a transformation $\mathrm{id}_{\mathsf{HA}} \to up' \cdot pf'$.

2.14 Definition. We call a logic (\mathcal{L}, ρ) for \mathcal{T} : Krip \rightarrow Pos *canonical* if there exists a natural transformation $\tau : pf \cdot \mathcal{L} \rightarrow \mathcal{T} \cdot pf'$ such that $\rho^{\flat} \circ \tau = id_{pf \cdot \mathcal{L}}$.

If (\mathcal{L}, ρ) is a canonical logic for \mathcal{T} then every (\mathcal{L}, j) -dialgebra $\mathcal{A} = (A, \alpha)$ gives rise to a \mathcal{T} -frame $\mathcal{A}_+ = (pf'A, \alpha_+)$, where α_+ is defined by

$$i(pf'A) = pf(jA) \xrightarrow{pf \alpha} pf(\mathcal{L}A) \xrightarrow{\tau_A} \mathcal{T}(pf'A).$$

If $h : \mathcal{A} \to \mathcal{B}$ is a morphism in $\text{Dialg}(\mathcal{L}, i)$, then it easily follows that pf'h is a (i, \mathcal{T}) -dialgebra morphism from \mathcal{B}_+ to \mathcal{A}_+ . In fact, the assignment $(\cdot)_+ : \text{Dialg}(\mathcal{L}, j) \to \text{Dialg}(i, \mathcal{T})$ defines a functor.

If θ : Prop $\to A$ is an assignment for an (\mathcal{L}, j) -dialgebra (A, α) then we can define a valuation V for the \mathcal{T} -frame $(pf'A, \alpha_+)$ by $V_{\theta}(p) = \{ \mathfrak{q} \in pf'A \mid p \in \mathfrak{q} \}$. This yields a map $(\cdot)_+$: AlgMod $(\mathcal{L}) \to Mod(\mathcal{T})$ given on objects by $(A, \alpha, \theta) \mapsto (pf'A, \alpha_+, V_{\theta})$. It is straightforward to verify that pf'h is a \mathcal{T} -model morphism from $(pf'A', \alpha'_+, V_{\theta'})$ to $(pf'A, \alpha_+, V_{\theta})$ whenever $h: (A, \alpha, \theta) \to (A', \alpha', \theta')$ is a morphism in AlgMod (\mathcal{L}) . So $(\cdot)_+$: AlgMod $(\mathcal{L}) \to Mod(\mathcal{T})$ defines a functor.

2.15 Definition. Let $\mathfrak{X} = (X, \gamma)$ be a \mathcal{T} -frame. Then the prime filter extension $\widehat{\mathfrak{X}}$ of \mathfrak{X} is defined to be $\widehat{\mathfrak{X}} = (\mathfrak{X}^+)_+$. Similarly, the prime filter extension of a \mathcal{T} -model \mathfrak{M} is $\widehat{\mathfrak{M}} = (\mathfrak{M}^+)_+$.

Let us give an explicit description of the prime filter extension of a \mathcal{T} -frame and \mathcal{T} -model. For a \mathcal{T} -frame \mathfrak{X} let $\hat{X} = pf'(up'X)$ be the collection of prime filters on X. Then the prime filter extension of \mathfrak{X} is of the form $(\mathfrak{X}^+)_+ = (\hat{X}, \hat{\gamma})$, where \hat{X} is ordered by inclusion, and $\hat{\gamma} : i\hat{X} \to \mathcal{T}\hat{X}$ is given by the composition

$$i(pf'(up'X)) \xrightarrow{pf(up(\mathcal{T}X))} pf(up(\mathcal{T}X)) \xrightarrow{pf\rho_X} pf(\mathcal{L}(up'X)) \xrightarrow{\tau_{up'X}} \mathcal{T}(pf'(up'X)).$$

The prime filter extension of a model (X, γ, V) is $(\widehat{X}, \widehat{\gamma}, \widehat{V})$ where

$$\widehat{V}(p) = \{ \mathfrak{q} \in \widehat{X} \mid V(p) \in \mathfrak{q} \}.$$

2.16 Example. If $\mathcal{T} = \mathcal{P}_{up}$ and the logic $(\mathcal{L}_{\Box}, \rho)$ is given as in Example 2.12 then we can find a right inverse $\tau : pf \cdot \mathcal{L}_{\Box} \to \mathcal{P}_{up} \cdot pf'$ of ρ^{\flat} . Define $\tau : pf \cdot \mathcal{L} \to \mathcal{P}_{up} \cdot pf' A$ by

$$\tau_A(\mathfrak{q}) \mapsto \bigcap \{ \widetilde{b} \mid \Box b \in \mathfrak{q} \}.$$

The proof that this is a well-defined natural transformation satisfying $\rho^{\flat} \circ \tau = \text{id}$ is similar to the proof of Lemma 4.6 in [17].

Let us explicitly compute the prime filter extension of a \Box -frame (X, γ) . Its underlying state space is the collection \widehat{X} of prime filters on X. Let $\mathfrak{q}, \mathfrak{p} \in \widehat{X}$ be prime filters. Then we have

$$\begin{split} \mathfrak{p} \in \widehat{\gamma}(\mathfrak{q}) \quad \text{iff} \quad \mathfrak{p} \in \bigcap \{ \widetilde{b} \mid \Box b \in \rho_X^{-1}((\gamma^{-1})^{-1})(\mathfrak{q}) \} \\ \quad \text{iff} \quad \Box b \in \rho_X^{-1}((\gamma^{-1})^{-1})(\mathfrak{q}) \text{ implies } b \in \mathfrak{p} \end{split}$$

The pre-condition can be simplified via

$$\begin{split} \Box b \in \rho_X^{-1}((\gamma^{-1})^{-1})(\mathfrak{q}) & \text{iff} \quad \{a \in up'X \mid a \subseteq b\} \in (\gamma^{-1})^{-1}(\mathfrak{q}) \\ & \text{iff} \quad \gamma^{-1}(\{a \in up'X \mid a \subseteq b\}) \in \mathfrak{q} \\ & \text{iff} \quad \{x \in X \mid \gamma(x) \subseteq b\} \in \mathfrak{q} \\ & \text{iff} \quad m_{\Box}(b) \in \mathfrak{q} \end{split}$$

Here $m_{\Box}(b) = \{x \in X \mid \gamma(x) \subseteq b\}$. Therefore $\widehat{\gamma} : \widehat{X} \to \mathcal{P}_{up}\widehat{X}$ is given by

$$\widehat{\gamma}(\mathfrak{q}) = \{ \mathfrak{p} \in pf'(up'X) \mid b \in \mathfrak{p} \text{ for all } b \in up'X \text{ with } m_{\Box}(b) \in \mathfrak{q} \}.$$

Jim de Groot

If we translate this back to a relation, we get that the prime filter extension of (X, R) is $(\widehat{X}, \widehat{R})$ where

$$\mathfrak{q}R\mathfrak{p}$$
 iff $m_{\Box}(b) \in \mathfrak{q}$ implies $b \in \mathfrak{p}$, for all $b \in up'X$.

This is in line with the definitions from intuitionistic modal logic from [37, Section 2], and also resembles the definition of ultrafilter extension for classical normal modal logic, see [6, Section 10.2].

Suppose that in addition (X, γ) is equipped with a valuation V. Then the prime filter extension of (X, γ, V) is given by $(\widehat{X}, \widehat{\gamma}, \widehat{V})$, where $\widehat{V}(p)$ consists of all prime filters in \widehat{X} that contain V(p).

2.17 Example. We claim that the logic $(\mathcal{L}_{\Delta}, \rho^{\Delta})$ from Example 2.13 is canonical. To avoid notational clutter, we ρ instead of ρ^{Δ} .

Let us first get an idea of the action of $\rho_A^{\flat} : \mathcal{H} \cdot pf'A \to pf \cdot \mathcal{L}_{\Delta}A$. Let $W \in \mathcal{H} \cdot pf'A$. A prime filter $Q \in pf(\mathcal{L}_{\Delta}A)$ is determined uniquely by the elements of the form Δa it contains, so we content ourselves with computing which of these are in $\rho_A^{\flat}(W)$. We have

$$\underline{\Delta}a \in \rho^{\flat}(W) \quad \text{iff} \quad \rho(\underline{\Delta}\widetilde{a}) \in \eta_{\mathcal{H} \cdot pf'A}(W)$$
$$\text{iff} \quad W \in \rho(\underline{\Delta}\widetilde{a})$$
$$\text{iff} \quad \widetilde{a} \in W$$

Recall that $\tilde{a} = \{ \mathfrak{q} \in pf'A \mid a \in \mathfrak{q} \}$. Based on this and inspired by the duality for descriptive monotone frames in [17, Section 8] we define $\tau_A : pf \cdot \mathcal{L}_{\Delta}A \to \mathcal{H} \cdot pf'A$ as follows. Let $Q \in pf \cdot \mathcal{L}_{\Delta}A$ and $D \in up'(pf'A)$, then

- If $D = \widetilde{a} = \{ \mathfrak{p} \in \widehat{X} \mid a \in \mathfrak{p} \}$ for some $a \in A$, then $\widetilde{a} \in \tau_A(Q)$ if $\underline{\Delta}a \in Q$;
- If D is closed, i.e., $D = \bigcap \{ \widetilde{a} \mid a \in A, D \subseteq \widetilde{a} \}$, then $D \in \tau_A(Q)$ if $D \subseteq \widetilde{a}$ implies $\Delta a \in Q$.
- For any other $D, D \in \tau_A(Q)$ if there is a closed $C \subseteq D$ such that $C \in \tau_A(Q)$.

We leave the (tedious) verification that τ is a well-defined natural transformation that is right inverse to ρ^{\flat} on components to the reader.

The prime filter extension $(\widehat{X}, \widehat{N})$ of a monotone frame (X, N) can now be given as follows. Let $\mathbf{q} \in \widehat{X}$ and $D \in up\widehat{X}$, then

- If $D = \tilde{a}$ for some $a \in A$, then $\tilde{a} \in \hat{N}(\mathfrak{q})$ iff $m_{\Delta}a \in \mathfrak{q}$, where $m_{\Delta}(a) = \{x \in X \mid a \in N(x)\};$
- If D is closed, then $D \in \widehat{N}(\mathfrak{q})$ iff $D \subseteq \widetilde{a}$ implies $\widetilde{a} \in \mathfrak{q}$.
- For any other $D, D \in \widehat{N}(\mathfrak{q})$ if there is a closed $C \subseteq D$ such that $C \in \widehat{N}(\mathfrak{q})$.

3 More modal intuitionistic logics

In this section we discuss four more examples of modal extensions of intuitionistic logic. Besides showcasing the versatility of the dialgebraic approach, these examples also add to a set of "atoms" from which we can construct new modal logics via the methods in Section 4. Each of these logics is canonical and given by predicate liftings and axioms. As such, they all satisfy the preconditions of the Goldblatt-Thomason theorems in Section 6.

We begin with two trivial logics, one for the case $\mathcal{T} = i$: Krip \rightarrow Pos and one for the constant functor \mathcal{C}_C to some finite poset C. Thereafter, we describe intuitionistic logic with

a diamond-like duality, similar to the framework for a \Box -modality from Example 2.7. Lastly, we describe the intuitionistic counterpart of neighbourhood semantics for a modal operator that satisfies no relations. (The classical treatment of such a modality can be found in e.g. [7, Chapter 7].)

3.1 A modality preserving distributive lattice operations

Consider $\mathcal{T} = i$: Krip \rightarrow Pos. As a frame, this can be viewed as a pair (X, t), where $t: X \rightarrow X$ is a monotone map on X. We construct a canonical logic for it.

A (trivial) predicate lifting for this functor is given by

$$\lambda_X = \mathrm{id}_X : \mathcal{U}p(iX) \to \mathcal{U}p(iX) : a \mapsto a.$$

It is easy to verify that the modality induced by this preserves finite meets and finite joins (i.e., all distributive lattice operations). Using these as equations, the procedure in Definition 2.10 simply yields the functor $j : HA \to DL$, and $\rho : j \cdot up' \to up \cdot i$ is the identity. As a consequence, ρ^{\flat} is also given by the identity. Therefore, setting $\tau : pf \cdot j \to i \cdot pf'$ as the identity shows that (j, id) is a canonical logic for Dialg(i, i).

3.2 A poset of nullary operators

For a finite poset C, let C_C : Krip \rightarrow Pos be the constant functor to C. That is, $C_C X = C$ for all $X \in \text{Krip}$ and $C_C f = \text{id}_C$ for all bounded morphisms f. We can define a logic (\mathcal{L}_C, ρ) for \mathcal{C}_C by choosing $\mathcal{L}_C : \text{HA} \rightarrow \text{DL}$ as the constant functor to the distributive lattice upC, and $\rho = \text{id}_{upC} : \mathcal{L}_C \cdot up' X \rightarrow up \cdot \mathcal{C}_C X$. Using the fact that up and pf restrict to a duality between finite posets and finite distributive lattices, one can verify that ρ^{\flat} is the identity on C. As a consequence, (\mathcal{L}_C, ρ) is canonical.

We can also obtain (\mathcal{L}_C, ρ) via predicate liftings and axioms. For each upset $a \in upC$ define a nullary predicate lifting

$$\lambda_X^a = a \in up \cdot \mathcal{C}_C X.$$

Let use abbreviate $\heartsuit^{\lambda^a} = \lceil a \rceil$ and $\underline{\heartsuit}^{\lambda^a} = \lceil a \rceil$. Together with the axioms

$$[\underline{C}] = \top, \qquad [\underline{\emptyset}] = \bot, \qquad [\underline{a}] \land [\underline{b}] = [\underline{a} \cap \underline{b}], \qquad [\underline{a}] \lor [\underline{b}] = [\underline{a} \cup \underline{b}],$$

where a and b range over upC, these give rise to the logic (\mathcal{L}_C, ρ) .

Indeed, many of the modal operators induced by these predicate liftings are definable from others. In particular, if C is ordered by the identity then we can do with $\lambda^c = \{c\} \in upC$ for all elements $c \in C$, and use the axioms

$$\underline{\lceil \{c\}\rceil} \land \underline{\lceil \{c'\}\rceil} = \bot \text{ if } c \neq c', \qquad \bigvee_{c \in C} \underline{\lceil \{c\}\rceil} = \top.$$

However, for the purpose of proving that (\mathcal{L}_C, ρ) is given by predicate liftings and relations, the first construction suffices.

3.3 A normal diamond modality

In [37], so-called \diamond -frames are used to interpret intuitionistic logic with an additional \diamond -like modality. We show that this is an instance of a dialgebraic intuitionistic logic.

3.1 Definition. A \diamond -frame is a tuple (X, R) consisting of a poset X and a relation R such that

$$(\geq_X \circ R \circ \geq_X) = R.$$

A \diamond -frame morphism from (X, R) to (X', R') is a function between the underlying sets that is bounded with respect to both the poset order as well as the relation R. We write WZ \diamond for the category of \diamond -frames and -morphisms.

The interpretation of $\Diamond \varphi$ in a (model based on a) \Diamond -frame (X, R) is as usual: a state x satisfies $\Diamond \varphi$ if it has an R-successor satisfying φ .

The category of \diamond -frames is isomorphic to a category of dialgebras. For a poset X define $\mathcal{P}_{dn}X$ as the collection of downwards closed subsets of X ordered by inclusion. If $f: X \to X'$ is a bounded morphism, define $\mathcal{P}_{dn}f: \mathcal{P}_{dn}X \to \mathcal{P}_{dn}X'$ by $\mathcal{P}_{dn}f(b) = \downarrow f[b] = \{x' \in X' \mid x' \leq_{X'} f(x) \text{ for some } x \in b\}$. Then $\mathcal{P}_{dn}: \text{Krip} \to \text{Pos}$ is a functor, and a straightforward verification shows:

3.2 Proposition. We have

$$WZ \diamondsuit \cong Dialg(i, \mathcal{P}_{dn}).$$

The modality \diamond can be described via the predicate lifting $\lambda^{\diamond} : \mathcal{U}p \cdot i \to \mathcal{U}p \cdot \mathcal{P}_{dn}$ defined as

$$\lambda_X^{\diamond}: \mathcal{U}p(iX) \to \mathcal{U}p(\mathcal{P}_{dn}X): a \mapsto \{b \in \mathcal{P}_{dn}X \mid a \cap b \neq \emptyset\}.$$

It is easy verify that the modality $\heartsuit^{\lambda^{\diamond}}$ induced by this coincides with \diamondsuit . Together with the axioms $\underline{\diamond} a \lor \underline{\diamond} b = \underline{\diamond} (a \lor b)$ and $\underline{\diamond} \bot = \bot$ we obtain a functor $\mathcal{L}_{\diamond} : \mathsf{HA} \to \mathsf{DL}$ that sends a Heyting algebra A to the free distributive lattice generated by $\{\underline{\diamond} a \mid a \in A\}$ modulo axioms stating that $\underline{\diamond}$ preserves finite joins. The corresponding natural transformation $\rho^{\diamond} : \mathcal{L}_{\diamond} \cdot up \to up \cdot \mathcal{P}_{dn}$ is given on generators by $\rho_X^{\diamond}(\underline{\diamond} a) = \{b \in \mathcal{P}_{dn} \mid a \cap b \neq \emptyset\}.$

To avoid notational clutter we temporarily write ρ rather than ρ^{\diamond} . Towards proving canonicity, let us compute the action of ρ^{\flat} on a downset D.

3.3 Lemma. Let A be a Heyting algebra, $D \in \mathcal{P}_{dn}(pf'A)$ and $a \in A$. Then we have

$$\underline{\diamond}a \in \rho_A^{\flat}(D) \quad iff \quad D \cap \widetilde{a} \neq \emptyset$$

Proof. We have

$$\begin{split} \underline{\diamond} a \in \rho_A^{\mathfrak{p}}(D) & \text{iff} \quad \underline{\diamond} a \in (\mathcal{L} \diamond \theta_A')^{-1}(\mathfrak{pf}\rho_{pf'A}(\eta(D))) & (\text{Definition of } \rho^{\mathfrak{p}}) \\ & \text{iff} \quad \underline{\diamond}(\theta_A'(a)) \in \mathfrak{pf}\rho_{pf'A}(\eta_{\mathcal{P}_{dn}},\mathfrak{pf'A}(D)) & (\text{Definition of } \mathcal{L}_{\diamond}) \\ & \text{iff} \quad \rho_{pf'A}(\underline{\diamond}(\theta_A'(a))) \in \eta_{\mathcal{P}_{dn}},\mathfrak{pf'A}(D) & (\text{Definition of } \mathfrak{pf}) \\ & \text{iff} \quad D \in \rho_{pf'A}(\underline{\diamond}\theta_A'(a)) & (\text{Definition of } \eta) \\ & \text{iff} \quad D \in \rho_{pf'A}(\underline{\diamond}\widetilde{a}) & (\text{Definition of } \theta) \\ & \text{iff} \quad D \cap \widetilde{a} \neq \emptyset & (\text{Definition of } \rho) \end{split}$$

as desired.

Inspired by this, we define $\tau : pf \cdot \mathcal{L}_{\diamond} \to \mathcal{P}_{dn} \cdot pf'$ on components by

$$\tau_A(Q) = pf'A \setminus \bigcup \{ \widetilde{a} \in up'A \mid \underline{\diamond}a \notin Q \}.$$
(1)

Clearly this is a monotone map, hence a morphism in Pos. We can prove that this is natural:

Jim de Groot

3.4 Lemma. The transformation $\tau : pf \cdot \mathcal{L}_{\diamond} \to \mathcal{P}_{dn} \cdot pf'$ is natural.

Proof. We need to show that for every Heyting algebra homomorphism $h: A \to A'$ the diagram

$$pf(\mathcal{L}_{\diamond}A) \xrightarrow{\tau_{A}} \mathcal{P}_{dn}(pf'A)$$
$$(\mathcal{L}_{\diamond}h)^{-1} \uparrow \qquad \uparrow \mathcal{P}_{dn}(h^{-1})$$
$$pf(\mathcal{L}_{\diamond}A') \xrightarrow{\tau_{A'}} \mathcal{P}_{dn}(pf'A')$$

commutes. To this end, let $Q' \in pf(\mathcal{L}_{\diamond}A')$ and $q \in pf'A$.

If $\mathbf{q} \in \mathcal{P}_{dn}(h^{-1})(\tau_{A'}(Q'))$ then there exists $\mathbf{p}' \in pf'A'$ such that $\mathbf{p}' \in \tau_{A'}(Q')$ and $\mathbf{q} \subseteq h^{-1}(\mathbf{p}')$. In order to prove that $\mathbf{q} \in \tau_A((\mathcal{L}_{\Diamond}h)^{-1}(Q'))$ we need to show that $\underline{\Diamond}a \notin (\mathcal{L}_{\Diamond}h)^{-1}(Q')$ implies $a \notin \mathbf{q}$. We have $\underline{\diamond}a \notin (\mathcal{L}_{\Diamond}h)^{-1}(Q')$ if and only if $\underline{\diamond}h(a) = \mathcal{L}_{\Diamond}h(\underline{\diamond}a) \notin Q'$. This in turn implies $h(a) \notin \mathbf{p}'$ and consequently $a \notin h^{-1}(\mathbf{p}')$. Since $\mathbf{q} \subseteq h^{-1}(\mathbf{p}')$ we find that $a \notin \mathbf{q}$.

Conversely, suppose $\mathfrak{q} \in \tau_A((\mathcal{L}_{\Diamond}h)^{-1}(Q'))$. To show that $\mathfrak{q} \in \mathcal{P}_{dn}(h^{-1})(\tau_{A'}(Q'))$ we need to construct a prime filter $\mathfrak{p}' \in pf'A'$ such that $\mathfrak{p}' \in \tau_{A'}(Q)$ and $\mathfrak{q} \subseteq h^{-1}(\mathfrak{p})$. We use the prime filter lemma. Let

$$I = \{ b' \in A' \mid \underline{\diamond} b' \notin Q' \}, \qquad F = \{ h(a) \in A' \mid a \in \mathfrak{q} \}.$$

Then the fact that $\underline{\diamond}$ preserves finite joins proves that I is an ideal in A'. Furthermore, $F \cap I = \emptyset$, since $h(a) \in F$ implies $a \in \mathfrak{q}$, which implies $\underline{\diamond} a \in (\mathcal{L}_{\diamond} h)^{-1}(A)$ and hence $\underline{\diamond} h(a) \in Q'$. Therefore $\uparrow F = \{c' \in A' \mid h(a) \leq c' \text{ for some } a \in \mathfrak{q}\}$ is a filter in A' disjoint from I. The prime filter lemma now yields a prime filter \mathfrak{p}' containing $\uparrow F$ disjoint from I. The fact that \mathfrak{p}' contains Fimplies $\mathfrak{q} \subseteq h^{-1}(\mathfrak{p}')$, and since \mathfrak{p}' is disjoint from I we have $\mathfrak{p}' \in \tau_{A'}(Q')$.

Finally, we prove canonicity of $(\mathcal{L}_{\diamond}, \rho)$.

3.5 Proposition. The logic $(\mathcal{L}_{\diamond}, \rho)$ is canonical for \mathcal{P}_{dn} .

Proof. We prove that the transformation τ is a right inverse of ρ^{\flat} , that is,

$$pf(\mathcal{L}_{\Diamond}A) \xrightarrow[\mathrm{id}]{\tau_A} \mathcal{P}_{dn}(pf'A) \xrightarrow[\mathrm{id}]{\rho_A'} pf(\mathcal{L}_{\Diamond}A)$$

commutes. Since prime filters $Q \in pf(\mathcal{L}_{\diamond}A)$ are determined uniquely by the elements of the form $\underline{\diamond}a$ they contain, where $a \in A$, it suffices to show that $\underline{\diamond}a \in \rho_A^{\flat} \circ \tau_A(Q)$ if and only if $\underline{\diamond}a \in Q$. We know from Lemma 3.3 that $\underline{\diamond}a \in \rho_A^{\flat}(\tau_A(Q))$ iff $\tau_A(Q) \cap \tilde{a} \neq \emptyset$, so we can get by with showing that

$$\tau_A(Q) \cap \widetilde{a} \neq \emptyset \quad \text{iff} \quad \underline{\diamond}a \in Q.$$

The direction from left to right holds by construction. For the converse direction we need to find some $q \in pf'A$ containing a such that $\underline{\diamond}b \notin Q$ implies $b \notin q$. This follows from a straightforward application of the prime filter lemma to the filter $\uparrow a = \{b \in A \mid a \leq b\}$ and the ideal $I = \{b \in A \mid \underline{\diamond}b \notin Q\}$.

Based on τ , we can describe the prime filter extensions of \diamond -frames. Let $\mathfrak{X} = (X, R)$ be a \diamond -frame, then $pf\mathfrak{X}$ has underlying poset \widehat{X} . The relation \widehat{R} is given by

$$\mathfrak{g} R\mathfrak{p}$$
 iff for all $a \in \mathfrak{u}\mathfrak{p}'X, a \in \mathfrak{p}$ implies $m_{\diamond}(a) \in \mathfrak{q}$

This corresponds with the notion of an ultrafilter extension for normal modal logic over a classical propositional base, see e.g. [2, Definition 2.57]. It is also in line with the definitions in [37, Section 2].

3.4 A free modality

Just as for classical propositional logic, we can augment intuitionistic logic with a unary modal operator that satisfies no axioms. The interpreting structure is a simple adaptation of neighbourhood frames [7]. We present these directly as dialgebras.

3.6 Definition. Define \mathcal{N} : Krip \rightarrow Pos on objects by sending a poset X to the set $\mathcal{U}pX$ ordered by inclusion. For a bounded morphism $f: X \rightarrow X'$ define $\mathcal{N}f: \mathcal{N}X \rightarrow \mathcal{N}X': W \mapsto \{a \in \mathcal{U}pX' \mid f^{-1}(a) \in W\}.$

An (intuitionistic) neighbourhood frame is an (i, \mathcal{N}) -dialgebra and morphisms between them are (i, \mathcal{N}) -dialgebra morphisms.

We can introduce a modal operator \odot via the predicate lifting

$$\lambda_X^{\odot}: \mathcal{U}p(iX) \to \mathcal{U}p(\mathcal{N}X): a \mapsto \{W \in \mathcal{N}X \mid a \in W\}.$$

This yields the logic $(\mathcal{L}_{\odot}, \rho)$, where \mathcal{L}_{\odot} is simply the composition of the forgetful functor $\mathsf{HA} \to \mathsf{Set}$ followed by the functor $\mathsf{Set} \to \mathsf{DL}$ taking a set to the free distributive lattice generated by it. That is, for a Heyting algebra $A, \mathcal{L}_{\odot}A$ is the free distributive lattice generated by $\{\underline{\odot}a \mid a \in A\}$. The natural transformation ρ is given by

$$\rho_X: \mathcal{L}_{\odot} \cdot up' X \to up \cdot \mathcal{N}X : \odot a \mapsto \{ W \in \mathcal{N}X \mid a \in W \}.$$

Now consider $\rho_A^{\flat} : \mathcal{N} \cdot pf'A \to pf \cdot \mathcal{L}_{\odot}A$. An easy computation shows that we have $\odot a \in \rho_A^{\flat}(W)$ iff $\tilde{a} \in W$ for all $a \in A$ and $W \in \mathcal{N} \cdot pf'A$. Therefore the transformation

$$\tau_A: pf \cdot \mathcal{L}_{\odot} A \to \mathcal{N} \cdot pf' A : Q \mapsto \{\widetilde{a} \mid \odot a \in Q\}$$

is clearly right inverse to ρ^{\flat} . A straightforward verification shows that moreover τ is natural. Therefore the logic induced by the predicate lifting λ^{\odot} and and empty set of axioms is canonical.

4 Combining modal logics

Given two functors $\mathcal{T}_1, \mathcal{T}_2$: Krip \to Pos, we can create a new functor in several ways. For example, we can define $\mathcal{T}_1 \times \mathcal{T}_2$ to send a poset X to $\mathcal{T}_1 X \times \mathcal{T}_2 X$ and a morphism $f: X \to X'$ in Krip to

$$(\mathcal{T}_1 \times \mathcal{T}_2)f : (\mathcal{T}_1 \times \mathcal{T}_2)X \to (\mathcal{T}_1 \times \mathcal{T}_2)X' : (t,s) \mapsto (\mathcal{T}_1f(t), \mathcal{T}_2f(s))$$

Similarly, taking pointwise coproducts gives rise to the functor $\mathcal{T}_1 + \mathcal{T}_2$: Krip \rightarrow Pos. Both of these are clearly functors. A third way of creating a new functor is a bit less common, and is more tailored towards ordered sets: We can define $(\mathcal{T}_1 \otimes \mathcal{T}_2)$ to send a poset X to the union of sets underlying $\mathcal{T}_1 X$ and $\mathcal{T}_2 X$, ordered by

$$x \le y \quad \text{iff} \quad \begin{cases} x, y \in \mathcal{T}_1 X \text{ and } x \le_1 y & \text{or} \\ x, y \in \mathcal{T}_2 X \text{ and } x \le_2 y & \text{or} \\ x \in \mathcal{T}_1 X \text{ and } y \in \mathcal{T}_2 X \end{cases}$$

In this section we show that each of these three constructions gives rise to a new functor $\operatorname{Krip} \to \operatorname{Pos}$ and we construct logics corresponding to these operations. That is, given logics (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) for functors \mathcal{T}_1 and \mathcal{T}_2 we construct a new logic for $\mathcal{T}_1 + \mathcal{T}_2$, $\mathcal{T}_1 \times \mathcal{T}_2$ and $\mathcal{T}_1 \otimes \mathcal{T}_2$. As an instance of such a constructed logic, we obtain $\Box \diamondsuit$ -frames and the corresponding logic from [37], see Example 4.2 below.

Moreover, we show that these constructions preserve canonicity and that the new logic is given by predicate liftings and axioms whenever (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are. Ultimately, this entails Theorem 4.18, which states that every intuitionistic polynomial functor has a canonical logic.

Throughout this section, we regularly omit the subscript of the natural transformations ρ , ρ^{\flat} and τ indicating the component, in an effort to avoid notational clutter. Whenever we do this, the component is clear from the context (the surrounding diagram).

4.1 Product functors and the coproduct of logics

We begin by investigating the product of functors \mathcal{T}_1 and \mathcal{T}_2 , and design a logic for $(\mathcal{T}_1 \times \mathcal{T}_2)$ -frames from logics for \mathcal{T}_1 - and \mathcal{T}_2 -frames.

4.1 Definition. Let $\mathcal{T}_1, \mathcal{T}_2 : \operatorname{Krip} \to \operatorname{Pos}$ be two functors and (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) two logics for them. Then $\mathcal{T}_1 \times \mathcal{T}_2 : \operatorname{Krip} \to \operatorname{Pos} : X \mapsto \mathcal{T}_1 X \times \mathcal{T}_2 X$ yields frame semantics for the sum of the logics (\mathcal{L}_i, ρ_i) . This sum (\mathcal{L}, ρ) is given as follows: $\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2$ sends a Heyting algebra A to the coproduct of $\mathcal{L}_1 A$ and $\mathcal{L}_2 A$ in DL, and ρ_X is obtained as the mediating map of the coproduct $\mathcal{L}_1 \cdot up' X + \mathcal{L}_2 \cdot up' X$ in the following diagram:

$$\begin{array}{cccc} \mathcal{L}_{1} \cdot up'X & \xrightarrow{\kappa_{1}} & \mathcal{L}_{1} \cdot up'X + \mathcal{L}_{2} \cdot up'X & \xleftarrow{\kappa_{2}} & \mathcal{L}_{1} \cdot up'X \\ & & \downarrow^{\rho_{1}} & & \downarrow^{\rho} & & \downarrow^{\rho_{2}} \\ up \cdot \mathcal{T}_{1}X & \xrightarrow{up\pi_{1}} & up(\mathcal{T}_{1}X \times \mathcal{T}_{2}X) & \xleftarrow{up\pi_{2}} & up \cdot \mathcal{T}_{1}X \end{array}$$

Here π_1, π_2 denote the projections $\mathcal{T}_1 X \times \mathcal{T}_2 X \to \mathcal{T}_i X$.

If both logics are given by predicate liftings and axioms, then so is their coproduct. Let $(\Lambda_i, \operatorname{Ax}_i)$ be sets of predicate liftings and axioms that gives rise the logic (\mathcal{L}_i, ρ_i) . For an *n*-ary predicate lifting $\lambda \in \Lambda_1$, define $\lambda^+ : up^n \to up \cdot (\mathcal{T}_1 \times \mathcal{T}_2)$ by

$$\lambda_X^+ : \mathcal{U}p^n(iX) \to \mathcal{U}p(\mathcal{T}_1X \times \mathcal{T}_2X) : a \mapsto \lambda_X(a) \times \mathcal{T}_2X.$$

Similarly define λ^* for predicate liftings $\lambda \in \Lambda_2$. Then $\Lambda = \{\lambda^+ \mid \lambda \in \Lambda_1\} \cup \{\lambda^* \mid \lambda \in \lambda_2\}$, together with the union $Ax_1 \cup Ax_2$ of both sets of axioms, gives rise to the logic (\mathcal{L}, ρ) defined abstractly above.

4.2 Example. Let $\mathcal{T}_1 = \mathcal{P}_{up}$ and $\mathcal{T}_2 = \mathcal{P}_{dn}$. Then $(\mathcal{P}_{up} \times \mathcal{P}_{dn})$ -frames correspond to the (full) \Box \diamond -frames from [37, Section 2]. We shall call these *WZ-frames*. Concretely, such frames are tuples of the form $(X, R_{\Box}, R_{\diamond})$, where X is a poset ordered by \leq_X and R_{\Box}, R_{\diamond} are binary relations on X that satisfy

$$(\leq_X \circ R_{\Box} \circ \leq_X) = R_{\Box}, \qquad (\geq_X \circ R_{\diamondsuit} \circ \geq_X) = R_{\diamondsuit}.$$

The coproduct $\mathcal{L}_{\Box} + \mathcal{L}_{\Diamond}$ of the logics described in Example 2.7 and Subsection 3.3 corresponds to the extension of intuitionistic logic with a box and a diamond satisfying

$$\Box \varphi \land \Box \psi \leftrightarrow \Box (\varphi \land \psi) \qquad \Box \top \leftrightarrow \top$$
$$\diamond \varphi \lor \diamond \psi \leftrightarrow \diamond (\varphi \lor \psi) \qquad \diamond \bot \leftrightarrow \bot$$

and no interaction axioms. This logic and its extensions are studied extensively by Wolter and Zakharyaschev [35, 36, 37]. We denote the logic by WZ. In Section 7 below we develop a Goldblatt-Thomason theorem for one of the extensions of WZ that was studied by Fischer Servi [10, 11] and Plotkin and Sterling [26].

Jim de Groot

4.3 Lemma. Let $\mathcal{T}_1, \mathcal{T}_2$: Krip \rightarrow Pos be two functors, with canonical logics (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) . Then the coproduct of these logics is a canonical logic for $\mathcal{T}_1 \times \mathcal{T}_2$.

Proof. We need to define a natural transformation $\tau : pf \cdot (\mathcal{L}_1 + \mathcal{L}_2) \to (\mathcal{T}_1 \times \mathcal{T}_2) \cdot pf'$ such that $\rho_A^{\flat} \circ \tau = \text{id.}$ By assumption we have τ_1, τ_2 such that $\rho_i^{\flat} \circ \tau_i = \text{id.}$ Define τ on components by setting τ_A to be the mediating map to the product $\mathcal{T}_1 \cdot pf'A \times \mathcal{T}_2 \cdot pf'A$ that makes

$$\begin{array}{cccc} \mathcal{T}_{1}(pf'A) & \xleftarrow{\pi_{1}} & \mathcal{T}_{1}(pf'A) \times \mathcal{T}_{2}(pf'A) & \xrightarrow{\pi_{2}} & \mathcal{T}_{2}(pf'A) \\ & \uparrow^{\tau_{1}} & & \uparrow^{\tau_{A}} & & \tau_{2} \uparrow \\ pf(\mathcal{L}_{1}A) & \xleftarrow{\sigma_{1}} & pf(\mathcal{L}_{1}A) \times pf(\mathcal{L}_{2}A) & \xrightarrow{\sigma_{2}} & pf(\mathcal{L}_{2}A) \end{array}$$

commute. Here the σ_i denote projection maps. Since $pf(\mathcal{L}_1A + \mathcal{L}_2A) \cong pf(\mathcal{L}_1A) \times pf(\mathcal{L}_2A)$ this is of the right type.

By definition of ρ and ρ^{\flat} , the upper two squares in the diagram below commute.

$$\begin{array}{c|c} pf(\mathcal{L}_{1}A) & \xleftarrow{\sigma_{1}} & pf(\mathcal{L}_{1}A) \times pf(\mathcal{L}_{2}A) & \xrightarrow{\sigma_{2}} & pf(\mathcal{L}_{2}A) \\ \hline \rho_{1}^{\flat} & & \rho_{2}^{\flat} & & \uparrow \rho_{2}^{\flat} \\ \hline \mathcal{T}_{1}(pf'A) & \xleftarrow{\pi_{1}} & \mathcal{T}_{1}(pf'A) \times \mathcal{T}_{2}(pf'A) & \xrightarrow{\pi_{2}} & \mathcal{T}_{2}(pf'A) \\ \hline \tau_{1} & & \uparrow \tau & & \uparrow \tau_{2} \\ pf(\mathcal{L}_{1}A) & \xleftarrow{\sigma_{1}} & pf(\mathcal{L}_{1}A) \times pf(\mathcal{L}_{2}A) & \xrightarrow{\sigma_{2}} & pf(\mathcal{L}_{2}A) \end{array}$$

Since the lower squares commute by definition, the outer shell commutes. Now since $\rho^{\flat} \circ \tau$ maps to a product, it is the unique map such that $\sigma_i \circ (\rho^{\flat} \circ \tau) = \rho_i^{\flat} \circ \tau_i \circ \sigma_i$ for i = 1, 2. As the identity on $pf \cdot \mathcal{L}_1A \times pf \cdot \mathcal{L}_2A$ also satisfies these conditions, we must have $\rho^{\flat} \circ \tau = id$.

4.2 A logic for the coproduct of functors

Our next goal is to construct a logic for $\mathcal{T}_1 + \mathcal{T}_2$. This resembles item 2 of Example 2.4 in [22]. Intuitively, the idea is to design a functor $\mathcal{L}_1 \star \mathcal{L}_2$ for the combination of the two logics for \mathcal{T}_1 and \mathcal{T}_2 in such a way that the prime filters of $(\mathcal{L}_1 \star \mathcal{L}_2)A$ correspond bijectively to $pf \mathcal{L}_1 A + pf \mathcal{L}_2 A$. We make use of the following auxiliary definition:

4.4 Definition. For distributive lattices A_1 and A_2 , define $A_1 \star A_2$ to be the free distributive lattice generated by $\phi_i a_i$, where i = 1, 2 and $a_i \in A_i$, modulo (the smallest congruence containing) the relations

$$\begin{split} & \Diamond_i \bot_i = \bot \\ & \Diamond_i a_i \land \Diamond_i b_i = \Diamond_i (a_i \land b_i) \\ & \Diamond_i a_i \lor \Diamond_i b_i = \Diamond_i (a_i \lor b_i) \\ & \Diamond_1 a_1 \land \Diamond_2 a_2 = \bot \\ & \Diamond_1 \top_1 \lor \Diamond_2 \top_2 = \top \end{split}$$

Intuitively, finite joins and binary meets of generators from the same A_i are computed as in A_i , binary meets of the form $\phi_1 a_1 \wedge \phi_2 a_2$ are killed, and joins of the form $\phi_1 a_1 \vee \phi_2 a_2$ are freely generated. As desired, prime filters of $A_1 \star A_2$ enjoy the following convenient property:

4.5 Lemma. We have $pf(A_1 \star A_2) \cong pfA_1 + pfA_2$.

Jim de Groot

Proof. A prime filter \mathfrak{p} of $A_1 \star A_2$ is determined uniquely by the generators of $A_1 \star A_2$ it contains. If \mathfrak{q} is a prime filter of A_i , then the set $\{ \diamond_i a_i \mid a_i \in q \} \cup \{ \top \}$ is a prime filter in $A_1 \star A_2$. This yields a map $pfA_1 + pfA_2 \rightarrow pf(A_1 \star A_2)$.

Conversely, suppose \mathfrak{p} is a prime filter of $A_1 \star A_2$. If \mathfrak{p} contains generators $\diamond a_1$ and $\diamond a_2$ with $a_i \in A_i$, then $\bot = \diamond_1 a_1 \land \diamond_2 a_2 \in \mathfrak{p}$, a contradiction. Moreover, the fact that $\diamond_1 \top_1 \lor \diamond_2 \top_2 = \top$ shows that either $\diamond_1 \top_1 \in \mathfrak{p}$ or $\diamond_2 \top_2 \in \mathfrak{p}$. Suppose $\diamond_i \top_i \in \mathfrak{p}$, then $\{a_i \in A_i \mid \diamond_i a_i \in p\}$ is a prime filter of A_i . This defines a map $pf(A_1 \star A_2) \rightarrow pfA_1 + pfA_2$.

It is easy to see that the two assignments are order-preserving and form a bijection. \Box

4.6 Definition. Let $\mathcal{T}_1, \mathcal{T}_2$: Krip \rightarrow Pos be functors with logics (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) . Define the functor $\mathcal{L}_1 \star \mathcal{L}_2$: HA \rightarrow DL by sending a Heyting algebra A to $\mathcal{L}_1 A \star \mathcal{L}_2 A$. For a homomorphism $h: A \rightarrow A'$ define $(\mathcal{L}_1 \star \mathcal{L}_2)h$ on generators by $(\mathcal{L}_1 \star \mathcal{L}_2)h(\phi_i a_i) = \phi_i(\mathcal{L}_i h(a_i))$, where *i* is either 1 or 2. It is straightforward to verify that this is a well-defined functor.

We define $\rho : (\mathcal{L}_1 \star \mathcal{L}_2) \cdot up' \to up \cdot (\mathcal{T}_1 + \mathcal{T}_2)$ on generators by

$$\rho_X(\diamond_i a_i) = \kappa_i [\rho_i(a_i)],$$

where $\kappa_i[-]$ denotes the direct image $up \cdot \mathcal{T}_i X \to up(\mathcal{T}_1 X + \mathcal{T}_2 X)$ of the inclusion $\kappa_i : \mathcal{T}_i X \to \mathcal{T}_1 X + \mathcal{T}_2 X$.

4.7 Lemma. The assignment $\rho : (\mathcal{L}_1 \star \mathcal{L}_2) \cdot up' \to up \cdot (\mathcal{T}_1 + \mathcal{T}_2)$ from Definition 4.6 is well defined and natural.

Proof. To show that ρ is well defined, we need to show that ρ preserves the equations from Definition 4.4. This follows from a straightforward verification, making use of the fact that we know precisely what $\mathcal{T}_1 X + \mathcal{T}_2 X$ looks like.

To see that ρ is natural we have to work a bit harder. Let $f: X \to X'$ be a morphism in Krip. Then we need to show that

commutes. It suffices to show that

$$\rho_X \circ (\mathcal{L}_1 \star \mathcal{L}_2) f(\diamond_i a'_i) = (\mathcal{T}_1 f + \mathcal{T}_2 f)^{-1} (\rho_{X'}(\diamond_i a'_i))$$

for every generator $\varphi_i a'_i$. To see that this is indeed the case, compute

$$\begin{split} \rho_X \circ ((\mathcal{L}_1 \star \mathcal{L}_2) f^{-1})(\phi_i a'_i) & \text{(Definition of } \mathcal{L}_1 \star \mathcal{L}_2) \\ &= \rho_X (\phi_i((\mathcal{L}_i f^{-1})(a'_i))) & \text{(Definition of } \mathcal{L}_1 \star \mathcal{L}_2) \\ &= \kappa_i [\rho_{i,X}(\mathcal{L}_i f^{-1})(a'_i)] & \text{(Definition of } \rho) \\ &= \kappa_i [(\mathcal{T}_i f)^{-1}(\rho_{i,X'}(a'_i))] & \text{(Naturality of } \rho_i) \\ &= (\mathcal{T}_1 f + \mathcal{T}_2 f)^{-1}(\kappa'_i [\rho_{i,X'}(a_i)]) & \text{(By Claim 1 below)} \\ &= (\mathcal{T}_1 f + \mathcal{T}_2 f)^{-1}(\rho_{X'}(\phi_{a_i})) & \text{(Definition of } \rho) \end{split}$$

Claim 1. Let $\kappa_i : \mathcal{T}_i X \to \mathcal{T}_1 X + \mathcal{T}_2 X$ and $\kappa'_i : \mathcal{T}_i X' \to \mathcal{T}_1 X' + \mathcal{T}_2 X'$ be inclusion maps. Then

$$up \cdot \mathcal{T}_{i}X \xrightarrow{\kappa_{i}[-]} up(\mathcal{T}_{1}X + \mathcal{T}_{2}X)$$
$$(\mathcal{T}_{i}f)^{-1} \uparrow \qquad \uparrow (\mathcal{T}_{1}f + \mathcal{T}_{2}f)^{-1}$$
$$up \cdot \mathcal{T}_{i}X' \xrightarrow{\kappa_{i}'[-]} up(\mathcal{T}_{1}X' + \mathcal{T}_{2}X')$$

commutes.

Proof of claim. This follows from a manual check: $\kappa'_i[-]$ sends an upset $a' \in up \cdot \mathcal{T}_i X'$ to a', viewed as a subset of $\mathcal{T}_1 X' + \mathcal{T}_2 X'$. Since $\mathcal{T}_1 f + \mathcal{T}_2 f$ maps elements from $\mathcal{T}_i X$ (viewed as elements of $\mathcal{T}_1 X + \mathcal{T}_2 X$) to elements in $\mathcal{T}_i X'$, we have $x \in (\mathcal{T}_1 f + \mathcal{T}_2 f)^{-1}(\kappa'_i[a'])$ if and only if $x \in (\mathcal{T}_i f)^{-1}(a')$. It follows that the diagram commutes.

This completes the proof of the lemma.

If (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are given by predicate liftings and equations, then so is $(\mathcal{L}_1 \star \mathcal{L}_2, \rho)$. For $\lambda \in \Lambda_i$, define

$$\lambda'_X: \mathcal{U}p^n(iX) \to \mathcal{U}p \cdot (\mathcal{T}_1 + \mathcal{T}_2)X: a \mapsto \kappa_i[\lambda_X(a)].$$

and let $\Lambda'_i = \{\lambda' \mid \lambda \in \Lambda_i\}$. Besides, define two nullary predicate liftings $\top_1 = \mathcal{T}_1 X \in up(\mathcal{T}_1 + \mathcal{T}_2)X$ and $\top_2 = \mathcal{T}_2 X \in up(\mathcal{T}_1 + \mathcal{T}_2)X$. Then the set of predicate liftings generating $(\mathcal{L}_1 \star \mathcal{L}_2, \rho)$ is

$$\Lambda = \Lambda'_1 \cup \Lambda'_2 \cup \{\top_1, \top_2\}.$$

The corresponding equations are

$$E = E'_1 \cup E'_2 \cup \{\underline{\lambda}_1 x_1 \land \underline{\lambda}_2 x_2 = \bot \mid \underline{\lambda}_i \in \Lambda'_i\} \cup \{\top_1 \lor \top_2 = \top\},\$$

where E'_i contains the equations from E_i but with every occurrence of $\underline{\lambda}$ replaced by $\underline{\lambda}'$ for $\lambda \in \Lambda_i$.

Intuitively, the resulting logic is a modal extension of intuitionistic logic with two sets of *incompatible* modalities. That is, a state cannot simultaneously satisfy modalities from both of these sets. The nullary predicate liftings \top_1 and \top_2 , given by $\top_i = \mathcal{T}_i X \in up(\mathcal{T}_1 + \mathcal{T}_2) X$ indicate which set of modalities can be potentially satisfied at a state. We call the logic $(\mathcal{L}_1 \star \mathcal{L}_2, \rho)$ the *incompatible union* of (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) . We know of no concrete examples of such a logic in the literature.

If (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are canonical, then so is their incompatible union.

4.8 Theorem. Suppose (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are canonical logics for \mathcal{T}_1 -frames and \mathcal{T}_2 -frames, respectively. Then the logic $(\mathcal{L}_1 \star \mathcal{L}_2, \rho)$ constructed in Definition 4.6 is canonical for $(\mathcal{T}_1 + \mathcal{T}_2)$ -frames.

Proof. We define a natural transformation $\tau : pf \cdot (\mathcal{L}_1 \star \mathcal{L}_2) \to (\mathcal{T}_1 + \mathcal{T}_2) \cdot pf'$ that satisfies $\rho^{\flat} \circ \tau = \text{id.}$ As a consequence of Lemma 4.5 we have $pf(\mathcal{L}_1 \star \mathcal{L}_2)A \cong pf(\mathcal{L}_1A) + pf(\mathcal{L}_2A)$ for $A \in \mathsf{HA}$. Therefore we can define τ_A as the mediating morphism making the following diagram commute:

Here the arrows called *in* are inclusions morphisms obtained from Lemma 4.5, and the κ_i are in inclusion maps of the coproduct $(\mathcal{T}_1 + \mathcal{T}_2) \cdot pf'A = \mathcal{T}_1 \cdot pf'A + \mathcal{T}_2 \cdot pf'A$. Naturality of τ follows from naturality of τ_1 and τ_2 .

In order to prove that $\rho^{\flat} \circ \tau = \text{id}$ it suffices to prove that the following diagram commutes:

$$\begin{array}{cccc} pf(\mathcal{L}_{1}A) & \stackrel{m_{1}}{\longrightarrow} pf(\mathcal{L}_{1}A) + pf(\mathcal{L}_{2}A) & \xleftarrow{m_{2}} pf(\mathcal{L}_{2}A) \\ & \downarrow^{\tau_{1}} & \downarrow^{\tau} & \tau_{2} \downarrow \\ & \downarrow^{\tau_{1}} & \downarrow^{\tau} & \mathcal{T}_{2} \cdot pf'A \\ & \downarrow^{\rho_{1}^{\flat}} & \downarrow^{\rho_{2}^{\flat}} & \mathcal{T}_{2} \cdot pf'A \\ & \downarrow^{\rho_{1}^{\flat}} & \downarrow^{\rho_{2}^{\flat}} & \rho_{2}^{\flat} \downarrow \\ & pf(\mathcal{L}_{1}A) \xrightarrow{i_{n_{1}}} pf(\mathcal{L}_{1}A) + pf(\mathcal{L}_{2}A) & \xleftarrow{i_{n_{2}}} pf(\mathcal{L}_{2}A) \end{array} \right)$$

Then $\rho_A^{\flat} \circ \tau_A$ is the mediating morphism arising from the coproduct $pf(\mathcal{L}_1A) + pf(\mathcal{L}_2A)$. Since the identity on $pf(\mathcal{L}_1A) + pf(\mathcal{L}_2A)$ also makes the diagram commute and the mediating map is unique, this implies $\rho^{\flat} \circ \tau = \text{id}$.

In order to prove that this diagram commutes, we unfold the definition of ρ^{\flat} . Consider the following diagram:

$$pf(\mathcal{L}_{1}A) \xrightarrow{in_{1}} pf(\mathcal{L}_{1}A) + pf(\mathcal{L}_{2}A) \xleftarrow{in_{2}} pf(\mathcal{L}_{2}A)$$

$$\downarrow^{\tau_{1}} (1) \downarrow^{\tau} (1) \tau_{2} \downarrow$$

$$\mathcal{T}_{1}(pf'A) \xrightarrow{\kappa_{1}} (\mathcal{T}_{1} + \mathcal{T}_{2})pf'A \xleftarrow{\kappa_{2}} \mathcal{T}_{2}(pf'A)$$

$$\downarrow^{\eta} (2) \downarrow^{\eta} (2) \downarrow^{\eta} (2) \downarrow^{\eta}$$

$$pf(up(\mathcal{T}_{1}(pf'A))) \xrightarrow{(\kappa_{1}^{-1})^{-1}} pf(up(\mathcal{T}_{1} + \mathcal{T}_{2})(pf'A)) \xleftarrow{(\kappa_{2}^{-1})^{-1}} pf(up(\mathcal{T}_{2}(pf'A)))$$

$$\downarrow^{pf}(\mathcal{L}_{1}(up'(pf'A))) \xrightarrow{\overline{in_{1}}} pf(\mathcal{L}_{1} \star \mathcal{L}_{2})(up'(pf'A)) \xleftarrow{\overline{in_{2}}} pf(\mathcal{L}_{2}(up'(pf'A)))$$

$$\downarrow^{pf \cdot \mathcal{L}_{1}\theta_{A}} (4) \downarrow^{pf \cdot \mathcal{L}_{1} \star \mathcal{L}_{2})(up'(pf'A)) \xleftarrow{in_{2}} pf(\mathcal{L}_{2}A)$$

The squares labelled (1) commute by definition of τ_A . Squares (2) commute by naturality of η . To see that the squares labelled (4) commute, first note that $pf \cdot (\mathcal{L}_1 \star \mathcal{L}_2) \cdot up' \cdot pf'A \cong pf \cdot \mathcal{L}_1 \cdot up' \cdot pf'A + pf \cdot \mathcal{L}_2 \cdot up' \cdot pf'A$. Then using the fact that in_i and in_i are given as in Lemma 4.5 it is straightforward to show that the squares commute.

Finally we prove that the squares labelled (3) commute. We give a proof for the left one. Let $Q \in pf \cdot up \cdot \mathcal{T}_1 \cdot pf'A$. In order to prove commutativity it suffices to show that for all generators $\phi_i a_i$ of $pf \cdot (\mathcal{L}_1 \star \mathcal{L}_2) \cdot up' \cdot pf'A$ we have

$$\diamond_i a_i \in \overline{in}_1(pf\rho_{1,pf'A}(Q)) \quad \text{iff} \quad \diamond_i a_i \in pf\rho_{pf'A}((\kappa_1^{-1})^{-1}(Q)) \tag{2}$$

Let us first consider generators of the form $\varphi_2 a_2$, where $a_2 \in \mathcal{L}_2 \cdot up' \cdot pf'A$. Clearly such a generator is never in the left hand side of (2). It is in the right hand side if and only if $\kappa_1^{-1}(\rho_{pf'A}(\varphi_2 a_2)) \in Q$, but $\rho_{pf'A}(\varphi_2 a_2) = \kappa_2[\rho_2(a_2)]$ and by the structure of the upsets of

 $(\mathcal{T}_1 + \mathcal{T}_2) \cdot pf'A$ we have $\kappa_1^{-1}(\kappa_2[\rho_2(a_2)]) = \emptyset$. Since Q is a filter it does not contain the empty set, so $\phi_2 a_2$ is not in the right hand side of (2) either. Now consider a generator of the form $\phi_1 a_1$. Then we have

$$\begin{split} & \phi_1 a_1 \in pf \, \rho_{pf'A}((\kappa_1^{-1})^{-1}(Q)) & \text{iff} \quad \rho_{pf'A}(\phi_1 a_1) \in (\kappa_1^{-1})^{-1}(Q) \\ & \text{iff} \quad \kappa_1[\rho_1(a_1)] \in (\kappa_1^{-1})^{-1}(Q) \\ & \text{iff} \quad \kappa_1^{-1}(\kappa_1[\rho_1(a_1)]) \in Q \\ & \text{iff} \quad \rho_1(a_1) \in Q \\ & \text{iff} \quad a_1 \in pf \, \rho_{1,pf'A}(Q) \\ & \text{iff} \quad \phi_1 a_1 \in \overline{in}_1(pf \, \rho_{1,pf'A}(Q)) \end{split}$$

In the fourth "iff" we use that κ_1 is injective, hence $\kappa_1^{-1} \circ \kappa_1$ is the identity.

This proves that the squares labelled (3) commute, and completes the proof of the theorem.

4.3 The ordered coproduct of functors

Contrary to the coproduct of posets X and Y, where we put them *in parellel*, we can also create a new poset by combining X and Y *in series*. That is, we take X and Y with their individual orders, and additionally stipulate every element of Y be greater than every element of X. This gives rise to the bifunctor \otimes : Pos \times Pos \rightarrow Pos defined as follows.

4.9 Definition. For posets X and Y define $X \otimes Y$ to be the poset with underlying set X + Y ordered by $\leq = \leq_X \cup \leq_Y \cup \{(x, y) \mid x \in X, y \in Y\}$. For morphisms $f : X \to X'$ and $g : Y \to Y'$ define $f \otimes g : X \otimes Y \to X' \otimes Y'$ by $(f \otimes g)(x) = f(x)$ for $x \in X$ and $(f \otimes g)(y) = g(y)$ for $y \in Y$. Then \otimes defines a bifunctor $\mathsf{Pos} \times \mathsf{Pos} \to \mathsf{Pos}$.

We call $X \otimes Y$ the ordered union of X and Y.

4.10 Remark. The element $X \otimes Y$ may alternatively be defined as the weighted colimit in Pos (viewed as a 2-category) of the diagram $X \xrightarrow{in_X} X + Y \xleftarrow{in_Y} Y$ with weight $\mathbf{1} \xrightarrow{0} \mathbf{2} \xleftarrow{1} \mathbf{1}$ where $0, 1 : \mathbf{1} \to \mathbf{2}$ denote the functions to $0, 1 \in \mathbf{2}, \mathbf{1}$ is the one-element poset and $\mathbf{2} = \{0, 1\}$ is a two-element poset with $0 \leq \mathbf{2}$ 1.

4.11 Definition. We write $\mathcal{T}_1 \otimes \mathcal{T}_2$: Krip \rightarrow Pos for the composition of \otimes with \mathcal{T}_1 and \mathcal{T}_2 .

Let (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) be logics for \mathcal{T}_1 and \mathcal{T}_2 . We combine these into a logic for $\mathcal{T}_1 \otimes \mathcal{T}_2$. Towards this, we first define a bifunctor \triangleright on DL. Intuitively, we define $A \triangleright B$ to be the distributive lattice one obtains from taking the disjoint union of A and B, and identifying the top element of B with the bottom element of A. As a consequence, every element of A is greater than every element of B.

4.12 Definition. For distributive lattices A and B, define $A \triangleright B$ to be the distributive lattice generated by the elements of A and B modulo the relations

$$\top = \top_A, \qquad \bot = \bot_B, \qquad \bot_A = \top_B,$$

and

$$a \wedge a' = a \wedge_A a', \qquad a \vee a' = a \vee_A a'$$

if both $a, a' \in A$, and similar for $b, b' \in B$.

As a consequence of the relations we then have $a \wedge b = b$ and $a \vee b = a$ for all $a \in A$ and $b \in B$. It can be checked that this yields a distributive lattice whose underlying set is simply the disjoint union of A and B with \perp_A and \top_B identified. Since \triangleright is given by generators and relations it gives rise to a bifunctor $\triangleright : DL \times DL \rightarrow DL$. Its action on morphisms $h : A \rightarrow A'$ and $g : B \rightarrow B'$ is given by

$$h \triangleright g : A \triangleright B \to A' \triangleright B' : c \mapsto \begin{cases} h(c) & \text{if } c \in A \\ g(c) & \text{if } c \in B \end{cases}$$

In particular, this is the unique map such that

$$\begin{array}{ccc} A & \xleftarrow{\xi_1} & A \triangleright B & \xrightarrow{\xi_2} & B \\ h & & & \downarrow h \triangleright g & & \downarrow g \\ A' & \xleftarrow{\xi_1'} & A' \triangleright B' & \xrightarrow{\xi_2'} & B' \end{array}$$

commutes, where ξ_1 denotes the quotient map that sends every element from B to \perp_A and ξ_2 is sends every $a \in A$ to \top_B , and similar for ξ'_1, ξ'_2 .

4.13 Lemma. There exists an isomorphism $up(X \otimes Y) \cong upX \triangleright upY$ that is natural in both X and Y.

Proof. The upsets of $X \otimes Y$ are precisely the upsets of Y, and sets of the form $a \cup X$ where $a \in upX$ and $a \neq \emptyset$. (The case $a = \emptyset$ yields the up-set X, which is already accounted for as an up-set of X.) It is easy to see that this bijection is in fact an isomorphism of distributive lattices. Naturality follows from inspecting the definitions of \otimes and \triangleright on morphisms.

We can then define the desired combination of \mathcal{L}_1 and \mathcal{L}_2 as $\mathcal{L}_1 \triangleright \mathcal{L}_2$.

4.14 Definition. Let (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) be logics for \mathcal{T}_1 and \mathcal{T}_2 . Define the logic (\mathcal{L}, ρ) for $\mathcal{T}_1 \otimes \mathcal{T}_2$ by $\mathcal{L} = \mathcal{L}_1 \triangleright \mathcal{L}_2$ and with ρ is given by the composition

$$\mathcal{L}_{1}(upX) \triangleright \mathcal{L}_{2}(upX) \xrightarrow{\rho_{1,X} \triangleright \rho_{2,X}} up(\mathcal{T}_{1}X) \triangleright up(\mathcal{T}_{2}X) \xrightarrow{\cong} up(\mathcal{T}_{1}X \otimes \mathcal{T}_{2}X)$$

Suppose both logics are given by predicate liftings and axioms, say, (\mathcal{L}_i, ρ_i) is given by $(\Lambda_i, \operatorname{Ax}_i)$. Then so is $(\mathcal{L}_1 \triangleright \mathcal{L}_2, \rho)$. Intuitively, we take the union of $(\Lambda_1, \operatorname{Ax}_1)$ and $(\Lambda_2, \operatorname{Ax}_2)$ and stipulate that every generator of the form $\underline{\heartsuit}^{\lambda_1}$ is bigger than every generator of the form $\underline{\heartsuit}^{\lambda_2}$, where $\lambda_1 \in \Lambda_1$ and $\lambda_2 \in \Lambda_2$. However, we do need one more additional nullary predicate lifting to take into account the element $\perp_A = \top_B$. We call this the separator predicate lifting, and denote it by \mathbb{T} . This is defined by $\mathbb{T}_X = \mathcal{T}_1 X \in up(\mathcal{T}_1 \otimes \mathcal{T}_2) X$. Let Ax be the union of $\operatorname{Ax}_1, \operatorname{Ax}_2$ and the collection

$$\underline{\bot} \leq \underline{\heartsuit}^{\lambda_1}(a_1, \dots, a_n), \qquad \underline{\heartsuit}^{\lambda_2}(a_1, \dots, a_n) \leq \underline{\bot}$$

where $\lambda_1 \in \Lambda_1, \lambda_2 \in \Lambda_2$. Then we have a collection of predicate liftings and axioms Ax whose induced logic is $(\mathcal{L}_1 \triangleright \mathcal{L}_2, \rho)$.

We complete this subsection by proving that $(\mathcal{L}_1 \triangleright \mathcal{L}_2, \rho)$ is canonical whenever (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are. We make use of the following lemma.

4.15 Lemma. We have an isomorphism $pf(A \triangleright B) \cong (pfA) \otimes (pfB)$ which is natural in A and B.

Jim de Groot

Proof. It is easy to see that we have a bijection between prime filters of $A \triangleright B$ and the disjoint union of pfA and pfB. Indeed, the prime filters of $A \triangleright B$ that do not contain $\bot_A = \top_B$ correspond bijectively with pfA, and prime filters of $A \triangleright B$ containing \top_B correspond bijectively to pfB. Furthermore, by construction of $A \triangleright B$ every prime filter of $A \triangleright B$ not containing \top_B is a subset of every prime filter of $A \triangleright B$ containing \top_B .

Naturality in both arguments can be verified by a straightforward computation.

4.16 Proposition. Suppose (\mathcal{L}_1, ρ_1) and (\mathcal{L}_2, ρ_2) are canonical logics for \mathcal{T}_1 -frames and \mathcal{T}_2 -frames, respectively. Then the logic $(\mathcal{L}_1 \triangleright \mathcal{L}_2, \rho)$ constructed in Definition 4.14 is canonical for $(\mathcal{T}_1 \otimes \mathcal{T}_2)$ -frames.

Proof. Define τ_A as the composition

$$pf(\mathcal{L}_1 \triangleright \mathcal{L}_2)A \xrightarrow{\cong} pf(\mathcal{L}_1A) \otimes pf(\mathcal{L}_2A) \xrightarrow{\tau_1 \otimes \tau_2} (\mathcal{T}_1 \otimes \mathcal{T}_2)(pf'A).$$

Then in particular the squares labelled (1) in the diagram below commute. Here the m_i and κ_i denote the obvious inclusion maps.

The argument to prove that $\tau_A \circ \rho_A^{\flat} = \operatorname{id}_{pf(\mathcal{L}_1 \triangleright \mathcal{L}_2)A}$ now proceeds as follows: Every prime filter $\mathbf{q} \in pf(\mathcal{L}_1 \triangleright \mathcal{L}_2)A$ is in the image of either ξ_1^{-1} or ξ_2^{-1} , let us say $\mathbf{q} = \xi_i^{-1}(\mathbf{q}')$. Moreover, the ξ_i^{-1} are injective. We then prove that the diagram commutes and as a consequence we have $\rho_A^{\flat}(\tau_A(\mathbf{q})) = \xi_i^{-1}(\rho_{i,A}^{\flat}(\tau_{i,A}(\mathbf{q}'))) = \xi_i^{-1}(\mathbf{q}') = \mathbf{q}$, since we assume the logics (\mathcal{L}_i, ρ_i) to be canonical.

So let us prove that all cells in the diagram commute. We have omitted subscripts of natural transformations in an effort to avoid notational clutter.

$$\begin{split} & \xi_{1}^{-1} & pf(\mathcal{L}_{1} \triangleright \mathcal{L}_{2})A & \xi_{2}^{-1} \\ & \downarrow \cong \\ & pf(\mathcal{L}_{1}A) & \xrightarrow{m_{1}} pf(\mathcal{L}_{1}A) \otimes pf(\mathcal{L}_{2}A) & \xleftarrow{m_{2}} pf(\mathcal{L}_{2}A) \\ & \tau_{1} & (1) & \downarrow \tau_{1} \otimes \tau_{2} & (1) & \downarrow \tau_{2} \\ & \mathcal{T}_{1}(pf'\mathcal{A}) & \xrightarrow{\kappa_{1}} \mathcal{T}_{1}(pf'\mathcal{A}) \otimes \mathcal{T}_{2}(pf'\mathcal{A}) & \xleftarrow{\kappa_{2}} \mathcal{T}_{2}(pf'\mathcal{A}) \\ & \downarrow & \eta & (2) & \downarrow & \eta & (2) & \downarrow & \eta \\ & & (\kappa_{1}^{-1})^{-1} & pf(up(\mathcal{T}_{1}(pf'\mathcal{A}) \otimes \mathcal{T}_{2}(pf'\mathcal{A})))) & (\kappa_{2}^{-1})^{-1} & \downarrow & \eta \\ & pf(up(\mathcal{T}_{1}(pf'\mathcal{A}))) & (3) & \cong & (3) & pf(up(\mathcal{T}_{2}(pf'\mathcal{A})))) \\ & pf(up(\mathcal{T}_{1}(pf'\mathcal{A}))) & (3) & \cong & (3) & pf(up(\mathcal{T}_{2}(pf'\mathcal{A})))) \\ & pf(\mathcal{L}_{1}(up'(pf'\mathcal{A}))) & (3) & \cong & (3) & pf(up(\mathcal{T}_{2}(pf'\mathcal{A})))) \\ & pf(\mathcal{L}_{1}(up'(pf'\mathcal{A}))) & (3) & \cong & (3) & pf(up(\mathcal{T}_{2}(pf'\mathcal{A})))) \\ & pf(\mathcal{L}_{1}(up'(pf'\mathcal{A}))) & (3) & \downarrow & pf(\rho_{1} \triangleright \rho_{2}) & (4) & \downarrow & pf(\rho_{2}) \\ & pf(\mathcal{L}_{1}\mathcal{A}) & (5) & \downarrow & pf(\mathcal{L}_{1} \land \mathcal{L}_{2}\mathcal{A}) & (5) & pf(\mathcal{L}_{2}(up'(pf'\mathcal{A})))) \\ & \downarrow & pf(\mathcal{L}_{1}\mathcal{A}) & (5) & \downarrow & pf(\mathcal{L}_{1}\mathcal{A} \triangleright \mathcal{L}_{2}\mathcal{A}) & (5) & pf(\mathcal{L}_{2}\mathcal{A}) \\ & pf(\mathcal{L}_{2}\mathcal{A}) & (5) & ($$

Commutativity of the top triangles follows from the proof of Lemma 4.15. Cells (2) commute by naturality of η . Commutativity of cells (3) follows from the proof of Lemma 4.13 Finally,

commutativity of the cells labelled (4) and (5) follows from applying pf to the defining diagram of the action of \triangleright on a pair of morphisms.

4.4 Intuitionistic Kripke polynomial functors

We now define a recursive class of functors $\mathcal{T} : \mathsf{Krip} \to \mathsf{Pos}$ that all have a canonical logic given by predicate liftings and axioms. This is an analog of (generalised) Kripke polynomial functors on Set, introduced in [29, Definition 1.1] and [22, Definition 2.2]. Therefore, we will refer to them as *intuitionistic polynomial functors*.

4.17 Definition. The class IPF of *intuitionistic polynomial functors* is defined recursively via

$$\mathbf{IPF} \ni \mathcal{T} ::= i \mid \mathcal{C}_C \mid \mathcal{P}_{up} \mid \mathcal{P}_{dn} \mid \mathcal{H} \mid \mathcal{N} \mid \mathcal{T} \times \mathcal{T} \mid \mathcal{T} + \mathcal{T} \mid \mathcal{T} \otimes \mathcal{T}$$

where C is a finite poset.

4.18 Theorem. Every functor $\mathcal{T} \in \mathbf{IPF}$ has a canonical logic given by predicate liftings and equations.

Proof. We prove this by induction on the structure of \mathcal{T} . The base cases we have seen in Examples 2.16 and 2.17 and in Section 3. The inductive steps follow from the preceding subsections.

5 Invariance

In this section we discuss preservation of validity of formulae. Ultimately, we prove that validity for formulae in a frame is preserved by regular subframes, p-morphic images, regular bisimilar images and disjoint unions, and that it is reflected by prime filter extensions. (That is, if a formula φ is valid on the prime filter extension $\hat{\mathfrak{X}}$ of some frame \mathfrak{X} , then it is also valid on \mathfrak{X} .)

Similarly, for models we prove that validity of formulae is preserved by p-morphic domains and images, m-bisimilar images and disjoint unions, and invariant under prime filter extensions.

Throughout this section, we assume given a functor $\mathcal{T} : \mathsf{Krip} \to \mathsf{Pos}$ and a logic for \mathcal{T} given by predicate liftings and axioms. Only when discussing prime filter extensions we require the logic to be canonical (to ensure existence of the prime filter extensions). We write $\mathfrak{X} = (X, \gamma)$ and $\mathfrak{X}' = (X', \gamma')$ for generic (i, \mathcal{T}) -dialgebras, and $\mathfrak{M} = (X, \gamma, V)$ and $\mathfrak{M}' = (X', \gamma', V')$ for generic \mathcal{T} -models.

5.1 Morphisms and generated subframes

The goal of this subsection is to prove that validity of a formula in a frame is preserved under regular subframes and p-morphic images, and validity of a formula in a model is preserved by p-morphic domains and images. We begin by showing that \mathcal{T} -model morphisms preserve the truth at individual states.

5.1 Proposition. Let $f : \mathfrak{M} \to \mathfrak{M}'$ be a \mathcal{T} -model morphism. Then for all $x \in X$ and $\varphi \in \mathbb{L}(\Lambda, \operatorname{Prop})$ we have

$$\mathfrak{M}, x \Vdash \varphi \quad iff \quad \mathfrak{M}', f(x) \Vdash \varphi.$$

Proof. Straightforward induction on the structure of φ .

Let $f : \mathfrak{M} \to \mathfrak{M}'$ be a \mathcal{T} -model morphism. Then we call \mathfrak{M} a *p*-morphic domain of \mathfrak{M}' . If f is surjective we say that \mathfrak{M}' is a *p*-morphic image of \mathfrak{M} . As a corollary of Proposition 5.1 we obtain:

5.2 Corollary. Let $f : \mathfrak{M} \to \mathfrak{M}'$ be a \mathcal{T} -model morphism. Then $\mathfrak{M}' \Vdash \varphi$ implies $\mathfrak{M} \Vdash \varphi$. If f is surjective then the converse also holds.

We are interested in the notion of a regular subframe. This plays that same rôle that a generated subframe plays in the Goldblatt-Thomason theorem of normal modal logics. We call these *regular* because they are frames that can be mapped into a frame via a dialgebra morphism whose underlying map is a regular monomorphism (which is precisely an embedding) in Krip.

5.3 Definition. Let $\mathfrak{X} = (X, \gamma)$ be a \mathcal{T} -frame. Then a regular subframe of (X, γ) is a \mathcal{T} -frame (X', γ') such that there exists an embedding $m : X' \to X$ in Krip such that $m : (X', \gamma') \to (X, \gamma)$ is a \mathcal{T} -frame morphism.

If \mathcal{T} preserves monos, then for every embedding $X' \to X$ there is at most one structure map $\gamma' : iX' \to \mathcal{T}X'$ such that $m : (X', \gamma') \to (X, \gamma)$ is a \mathcal{T} -frame morphism. Although this is often the case, we do not need it in the proof of the Goldblatt-Thomason theorem below, so we shall not assume such a property.

5.4 Example. Let us revisit the example of \Box -frames (i.e., \mathcal{P}_{up} -frames, or (i, \mathcal{P}_{up}) -dialgebras). Let (X, R) be a \Box -frame and $X' \subseteq X$. Then (X', R') is a regular subframe of (X, R) if

- for all $x, y \in X'$, xR'y iff xRy;
- for all $x \in X'$ and $z \in X$, if xRz then $z \in X'$.

A straightforward verification shows that this coincides precisely with the notion of a regular subframe for \mathcal{P}_{uv} -frames from the definition above.

A similar description goes for \diamond -frames, except that the second bullet point should be replaced by: for all $x \in X'$ and $z \in X$, if xRz then there exists $z' \in X'$ such that xR'z' and $z \leq_X z'$.

5.5 Example. We can similarly describe regular subframes of monotone frames. A regular subframe of a monotone frame (X, N) is a monotone frame (X', N') such that X' is (isomorphic to) a sub-poset of X and inherits its order from X, and satisfies for all $x' \in X'$ and $a \subseteq X$:

$$a \in N(x)$$
 iff $a \cap X' \in N'(x)$.

5.6 Proposition. Let \mathfrak{X} be a generated subframe of \mathfrak{X}' . Then $\mathfrak{X}' \Vdash \varphi$ implies $\mathfrak{X} \Vdash \varphi$.

Proof. Suppose V is a valuation for \mathfrak{X} . Define $V' : \operatorname{Prop} \to \mathcal{U}pX'$ by $V'(p) = \uparrow (f[V(p)])$. Then by construction f is a \mathcal{T} -model morphism from (\mathfrak{X}, V) to (\mathfrak{X}', V') and the statement follows from Corollary 5.2.

5.7 Definition. Let $\mathfrak{X} = (X, \gamma)$ be a \mathcal{T} -frame. Then a \mathcal{T} -frame $\mathfrak{Z} = (Z, z)$ is a *p*-morphic image of \mathfrak{X} if there exists a surjective \mathcal{T} -frame morphism $f : \mathfrak{X} \to \mathfrak{Z}$.

5.8 Proposition. If \mathfrak{Z} is a p-morphic image of \mathfrak{X} then $\mathfrak{X} \Vdash \varphi$ implies $\mathfrak{Z} \Vdash \varphi$.

Proof. Let V' be a valuation for \mathfrak{X}' . Then setting $V = \mathcal{U}pf \circ V'$ yields a valuation for \mathfrak{X} and the statement follows from Corollary 5.2.

5.2 Dialgebra Bisimulations

Next we consider the notion of a *dialgebra bisimulation* between \mathcal{T} -frames and \mathcal{T} -models. We show that a class of \mathcal{T} -frames is closed under so-called regular bisimilar images if and only if it is closed under regular subframes and p-morphic images. A class of \mathcal{T} -models is closed under m-bisimilar images if and only if it is closed under p-morphic domains and images.

5.9 Definition. Let $\mathfrak{X}_1 = (X_1, \gamma_1)$ and $\mathfrak{X}_2 = (X_2, \gamma_2)$ be \mathcal{T} -frames. A sub-poset $B \subseteq X_1 \times X_2$ is called a *(dialgebra) bisimulation* if the projections $\pi_i : B \to X_i$ are bounded morphisms and there exists a monotone map $\beta : iB \to \mathcal{T}B$ making

$$\begin{array}{cccc} iX_1 & \stackrel{i\pi_1}{\longleftarrow} & iB \xrightarrow{i\pi_2} & iX_2 \\ \gamma_1 & & & & & & & \\ \gamma_1 & & & & & & & \\ \mathcal{T}X_1 & \stackrel{\mathcal{T}\pi_1}{\longleftarrow} & \mathcal{T}B \xrightarrow{\mathcal{T}\pi_2} & \mathcal{T}X_2 \end{array}$$

commute. A model bisimulation or m-bisimulation between \mathcal{T} -models (X_1, γ_1, V_1) and (X_2, γ_2, V_2) is a ρ -bisimulation B between the underlying \mathcal{T} -frames that additionally satisfies

$$x_1 \in V_1(p)$$
 iff $x_2 \in V_2(p)$

for all $(x_1, x_2) \in B$ and $p \in \text{Prop.}$

5.10 Example. Let $\mathfrak{X}_1 = (X_1, R_1)$ and $\mathfrak{X}_2 = (X_2, R_2)$ be two \Box -frames. Then a sub-poset $B \subseteq X_1 \times X_2$ is a dialgebra bisimulation between \mathfrak{X}_1 and \mathfrak{X}_2 if for all $(x_1, x_2) \in B$ the following conditions hold:

- If $x_1 \leq_1 y_1$ then there exists $y_2 \in X_2$ such that $x_2 \leq_2 y_2$ and $(y_1, y_2) \in B$;
- If $x_2 \leq y_2$ then there exists $y_1 \in X_1$ such that $x_1 \leq y_1$ and $(y_1, y_2) \in B$;
- If $x_1R_1y_1$ then there exists $y_2 \in X_2$ such that $x_2R_2y_2$ and $(y_1, y_2) \in B$;
- If $x_2R_2y_2$ then there exists $y_1 \in X_1$ such that $x_1R_1y_1$ and $(y_1, y_2) \in B$.

One gets the definition of a bisimulation between \Box -models by adding $x_1 \in V_1(p)$ iff $x_2 \in V_2(p)$. A Hennessy-Milner theorem for this notion of bisimulation has been given in [16].

Bisimulations between \diamond -frames can be described similarly.

5.11 Example. Let $\mathfrak{X}_1 = (X_1, N_1)$ and $\mathfrak{X}_2 = (X_2, N_2)$ be two monotone frames. Then a sub-poset $B \subseteq X_1 \times X_2$ is a dialgebra bisimulation between \mathfrak{X}_1 and \mathfrak{X}_2 for for all $(x_1, x_2) \in B$ we have:

- If $x_1 \leq_1 y_1$ then there exists $y_2 \in X_2$ such that $x_2 \leq_2 y_2$ and $(y_1, y_2) \in B$;
- If $x_2 \leq y_2$ then there exists $y_1 \in X_1$ such that $x_1 \leq y_1$ and $(y_1, y_2) \in B$;
- If $a_1 \in N_1(x_1)$ then there exists $a_2 \in N_2(x_2)$ such that $a_2 \subseteq B[a_1]$ and $B^{-1}[a_2] \cap a_1 \in N_1(x_1)$;
- If $a_2 \in N_2(x_2)$ then there exists $a_1 \in N_1(x_1)$ such that $a_1 \subseteq B^{-1}[a_2]$ and $B[a_1] \cap a_2 \in N_2(x_2)$;.

Here $B[a_1] = \{y_2 \in X_2 \mid (y_1, y_2) \in B \text{ for some } y_1 \in a_1\}$, and we similarly define $B^{-1}[a_2]$. The proof that this corresponds exactly to the notion of a dialgebra bisimulation is similar to the proof of [18, Proposition 8.23].

5.12 Proposition. Let B be an m-bisimulation between \mathcal{T} -models \mathfrak{M}_1 and \mathfrak{M}_2 . Then $(x_1, x_2) \in B$ implies

$$\mathfrak{M}_1, x_1 \Vdash \varphi \quad iff \quad \mathfrak{M}_2, x_2 \Vdash \varphi$$

for all $\mathbb{L}(\Lambda)$ -formulae φ .

Proof. Let β be the map $iB \to \mathcal{T}B$ we get from the definition of a bisimulation and define

 $V_B : \operatorname{Prop} \to B : p \mapsto \{(y_1, y_2) \in B \mid y_1 \in V_1(p), y_2 \in V_2(p)\}.$

Then it is easy to see that $\mathfrak{B} = (B, \beta, V_B)$ is a \mathcal{T} -model and the projections π_i are \mathcal{T} -model morphisms from \mathfrak{B} to \mathfrak{M}_i , so the result follows form Proposition 5.1.

5.13 Definition. Call a \mathcal{T} -model $\mathfrak{M}_2 = (X_2, \gamma_2, V_2)$ a *m*-bisimilar image of a \mathcal{T} -model \mathfrak{M}_1 if there exists a bisimulation B between them such that $\pi_2 : B \to X_2$ is surjective.

5.14 Proposition. Let \mathfrak{M}_2 be a m-bisimilar image of \mathfrak{M}_1 . Then $\mathfrak{M}_1 \Vdash \varphi$ implies $\mathfrak{M}_2 \Vdash \varphi$.

Proof. If $\mathfrak{M}_1 \Vdash \varphi$ then $\mathfrak{M}_1, x_1 \Vdash \varphi$ for all states x_1 in \mathfrak{M}_1 . Since \mathfrak{M}_2 is a bisimilar image of \mathfrak{M}_1 , for each state $x_2 \in \mathfrak{M}_2$ we can find $x_1 \in \mathfrak{M}_1$ such that $(x_1, x_2) \in B$. As a consequence of Proposition 5.12 this entails $\mathfrak{M}_2, x_2 \Vdash \varphi$ for all states of \mathfrak{M}_2 , hence $\mathfrak{M}_2 \Vdash \varphi$.

As for morphisms, a counterpart for this preservation result for frames requires a slightly stronger notion of bisimilar image; we need the notion of a *regular* bisimilar image. This has an additional requirement that π_1 is an embedding, which allows us to translate valuations for the image into valuations for the frame we start with and thus prove a preservation result.

5.15 Definition. Let $\mathfrak{X}_1 = (X_1, \gamma_1)$ and $\mathfrak{X}_2 = (X_2, \gamma_2)$ be two \mathcal{T} -frames. We call \mathfrak{X}_2 a regular bisimilar images of \mathfrak{X}_1 if there exists a bisimulation B between them such that $\pi_1 : B \to X_1$ is an embedding and $\pi_2 : B \to X_2$ is surjective.

5.16 Proposition. Let \mathfrak{X}_2 be a regular bisimilar image of \mathfrak{X}_1 . Then $\mathfrak{X}_1 \Vdash \varphi$ implies $\mathfrak{X}_2 \Vdash \varphi$.

Proof. This follows immediately from Propositions 5.6 and 5.8.

Finally, we describe when a class of \mathcal{T} -frames is closed under regular bisimilar images in terms of closure under generated subframes and p-morphic images, and similar for \mathcal{T} -models. We make use of the following Lemma, the proof of which is straightforward.

5.17 Lemma. 1. The graph of a \mathcal{T} -frame morphism is a bisimulation.

2. The graph of a \mathcal{T} -model morphism is an m-bisimulation.

5.18 Proposition. A class $\mathcal{K} \subseteq \text{Dialg}(i, \mathcal{T})$ of \mathcal{T} -frames is closed under regular bisimilar images if and only if it is closed under p-morphic images and regular subframes.

Proof. The direction from left to right follows from Lemma 5.17. Conversely, suppose \mathfrak{X}_2 is the regular bisimilar image of \mathfrak{X}_1 and let the bisimulation B be a witness of that. Then there exists $\beta : iB \to \mathcal{T}B$ such that π_1 and π_2 are \mathcal{T} -frame morphisms. Since π_1 is an embedding and \mathcal{K} is closed under generated subframes we have $(B, \beta) \in \mathcal{K}$. Closure of \mathcal{K} under p-morphic images then entails $\mathfrak{X}_2 \in \mathcal{K}$.

Jim de Groot

5.19 Proposition. A class $\mathcal{M} \subseteq \mathsf{Mod}(\mathcal{T})$ of \mathcal{T} -models is closed under m-bisimilar images if and only if it is closed under p-morphic images and domains.

Proof. The direction from left to right follows from Lemma 5.17. Conversely, let $\mathfrak{M}_2 = (X_1, \gamma_1, V_1)$ be the m-bisimilar image of $\mathfrak{M}_1 = (X_2, \gamma_2, V_2)$ and define $\mathfrak{B} = (B, \beta, V_B)$ as in the proof of Proposition 5.12. The result then follows from the fact that \mathfrak{B} is a p-morphic domain of \mathfrak{M}_1 and \mathfrak{M}_2 is a p-morphic image of \mathfrak{B} .

5.3 Disjoint unions

Both Pos and Krip have all small colimits and small colimits in Krip are computed as in Pos (see Appendix A.1). The coproduct of a family X_i of objects in either category is given by the disjoint union $X = \coprod X_i$ of the X_i ordered by $x \leq x'$ iff $x, x' \in X_i$ for the same *i* and $x \leq_i x'$. Here \leq_i denotes the order on X_i .

As a consequence of the fact that Krip has all small coproducts, the category $\text{Dialg}(i, \mathcal{T})$ has all small coproducts as well. Let (X_j, γ_j) be a family of (i, \mathcal{T}) -dialgebras. Then their coproduct or disjoint union $\coprod (X_j, \gamma_j)$ has state-space $\coprod X_i$ (computed in Krip). As a consequence of the observation above, we have $i(\coprod X_i) = \coprod (iX_j)$. We can then obtain the structure map for $\coprod (X_j, \gamma_j)$ as the unique mediating map γ in Pos that satisfies

$$\begin{array}{c} iX_k \xrightarrow{i\kappa_k} & \coprod iX_j \\ \gamma_k \downarrow & \downarrow \gamma \\ \mathcal{T}X_k \xrightarrow{\mathcal{T}\kappa_k} & \mathcal{T}(\coprod X_j) \end{array}$$

for all k. Here $\kappa_k : X_k \to \prod X_k$ denotes the inclusion map (in Krip).

This construction extends to models. If (X_j, γ_j, V_j) is a family of \mathcal{T} -models, then we can define their *disjoint union* or coproduct as (X, γ, V) where $(X, \gamma) = \coprod (X_j, \gamma_j)$ and V is defined by

$$V(p) = \bigcup V_j(p).$$

5.20 Example. Consider \Box -frames. If we translate all this, then the disjoint union of \Box -frames (X_j, R_j) is given by (X, R), where X is the disjoint union in Krip of the X_j and xRy iff $x, y \in X_j$ for some j and xR_jy .

5.21 Example. The product of two monotone frames (X_1, N_1) and (X_2, N_2) is the monotone frame (X, N), where X is the coproduct of X_1 and X_2 in Krip and N is defined by

$$a \in N(x_i)$$
 iff $a \cap X_i \in N_i(x_i)$

where $x_i \in X_i$ and $a \subseteq X$.

5.22 Proposition. Suppose $(X_j, \gamma_j, V_j) \Vdash \varphi$, then $\coprod (X_j, \gamma_j, V_j) \Vdash \varphi$.

Proof. This follows from the fact that each element x in the state-space of $\coprod (X_j, \gamma_j, V_j)$ lies in the image of one of the inclusion morphisms, i.e., $x = \kappa_k(x_k)$ for some $k \in J$ and $x_k \in X_k$. By assumption $x_k \Vdash \varphi$, so by Proposition 5.1 $x = \kappa_k(x_k) \Vdash \varphi$.

5.23 Proposition. Let $\mathfrak{X}_j = (X_j, \gamma_j)$ be a family of (i, \mathcal{T}) -dialgebras indexed by some set J and assume $\mathfrak{X}_j \Vdash \varphi$ for all $j \in J$. Then $\mathfrak{X} = \coprod \mathfrak{X}_j \Vdash \varphi$.

Jim de Groot

Proof. Let V be any valuation for \mathfrak{X} . Then setting

$$V_i: \operatorname{Prop} \to \mathcal{U}pX_i: p \mapsto \{x \in X_i \mid \kappa_i(x) \in V(p)\}$$

yields a valuation for each \mathfrak{X}_j that makes the inclusion map κ_j a \mathcal{T} -model morphism. Since $\mathfrak{X}_j \Vdash \varphi$ we have $(\mathfrak{X}_j, V_j) \Vdash \varphi$. It then follows from Proposition 5.22 that $(\mathfrak{X}, V) \Vdash \varphi$. Since V was chosen arbitrarily this implies $\mathfrak{X} \Vdash \varphi$.

5.4 Prime filter extensions

In this subsection we work with the following assumption:

5.24 Assumption. Assume given a functor \mathcal{T} : Krip \rightarrow Pos and a logic (\mathcal{L}, ρ) for \mathcal{T} -frames that is *canonical* and given by predicate liftings and axioms.

We have seen that this implies the existence of a right inverse τ of ρ^{\flat} , which in turn implies the existence of a notion of *prime filter extensions*. In this subsection we prove that this notion of prime filter extension satisfies:

- If \mathfrak{X} is a \mathcal{T} -frame and $\widehat{\mathfrak{X}} \Vdash \varphi$ then $\mathfrak{X} \Vdash \varphi$;
- If \mathfrak{M} is a \mathcal{T} -model then $\mathfrak{M} \Vdash \varphi$ iff $\widehat{\mathfrak{M}} \Vdash \varphi$.

Both of these are corollaries of the following proposition.

5.25 Proposition. For all $\varphi \in \mathbb{L}(\Lambda)$ and $\mathfrak{q} \in \widehat{\mathfrak{M}}$ we have

$$\widehat{\mathfrak{M}}, \mathfrak{q} \Vdash \varphi \quad iff \quad \llbracket \varphi \rrbracket^{\mathfrak{M}} \in \mathfrak{q}.$$

Proof. We prove the statement by induction on the structure of φ . If φ is a proposition letter then by definition of \hat{V} we have

$$\mathfrak{q} \Vdash p \quad \text{iff} \quad \mathfrak{q} \in \widehat{V}(p) \quad \text{iff} \quad \llbracket p \rrbracket^{\mathfrak{M}} = V(p) \in \mathfrak{q}.$$

The inductive cases for \bot, \top, \land and \lor are straightforward.

Consider the case $\varphi = \varphi_1 \to \varphi_2$ and suppose $\llbracket \varphi_1 \to \varphi_2 \rrbracket^{\mathfrak{M}} \in \mathfrak{q}$. If $\mathfrak{q} \subseteq \mathfrak{p}$ and $\mathfrak{p} \Vdash \varphi_1$, then by the inductive hypothesis we have $\llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \in \mathfrak{p}$ and since \mathfrak{p} is a filter also $\llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \cap \llbracket \varphi_1 \to \varphi_2 \rrbracket^{\mathfrak{M}} \in \mathfrak{p}$. By definition of the interpretation of formulae we have $\llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \cap \llbracket \varphi_1 \to \varphi_2 \rrbracket^{\mathfrak{M}} \subseteq \llbracket \varphi_2 \rrbracket^{\mathfrak{M}}$ and therefore $\llbracket \varphi_2 \rrbracket^{\mathfrak{M}} \in \mathfrak{p}$. It then follows again from the induction hypothesis that $\mathfrak{p} \Vdash \varphi_2$. This proves that $\mathfrak{q} \Vdash \varphi_1 \to \varphi_2$.

Conversely, assume $\mathbf{q} \Vdash \varphi_1 \to \varphi_2$, and suppose towards a contradiction that there is no $a \in \mathbf{q}$ such that $a \cap \llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \subseteq \llbracket \varphi_2 \rrbracket^{\mathfrak{M}}$. Then $\llbracket \varphi_2 \rrbracket^{\mathfrak{M}}$ is not in the filter generated by \mathbf{q} and $\{\llbracket \varphi_1 \rrbracket^{\mathfrak{M}}\}$, so there exists a filter \mathbf{p} such that $\mathbf{q} \subseteq \mathbf{p}$ and $\llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \in \mathbf{p}$ and $\llbracket \varphi_2 \rrbracket^{\mathfrak{M}} \notin \mathbf{p}$. But then $\mathbf{q} \nvDash \varphi_1 \to \varphi_2$, a contradiction. So we have $a \in \mathbf{q}$ such that $a \cap \llbracket \varphi_1 \rrbracket^{\mathfrak{M}} \subseteq \llbracket \varphi_2 \rrbracket^{\mathfrak{M}}$. But this implies $a \subseteq \llbracket \varphi_1 \to \varphi_2 \rrbracket^{\mathfrak{M}}$, and since \mathbf{q} is a filter we find $\llbracket \varphi_1 \to \varphi_2 \rrbracket^{\mathfrak{M}} \in \mathbf{q}$.

Finally, for the modal case we take a slight detour. It follows from the proof of [17, Proposition 6.10] that the unit $\theta'_{up'X} : up'X \to up'(pf'(up'X))$ is a (\mathcal{L}, j) -dialgebra morphism from \mathfrak{X}^+ to $(\hat{\mathfrak{X}})^+$, i.e., the following diagram commutes:

Jim de Groot

Therefore we have

$$\begin{split} \llbracket \heartsuit^{\lambda}(\varphi) \rrbracket^{pf\mathfrak{M}} &= \widehat{\gamma}^{*}(\underbrace{\heartsuit^{\lambda}}(\llbracket \varphi \rrbracket^{pf\mathfrak{M}})) \\ &= \widehat{\gamma}^{*}(\underbrace{\heartsuit^{\lambda}}(\llbracket \varphi \rrbracket^{\mathfrak{M}})) \qquad \text{(By induction)} \\ &= \widehat{\gamma}^{*}(\mathcal{L}\theta'_{upX}(\underbrace{\heartsuit^{\lambda}}(\llbracket \varphi \rrbracket^{\mathfrak{M}}))) \qquad \text{(Definition of } \zeta) \\ &= \theta'_{upX}(\gamma^{*}(\underbrace{\heartsuit^{\lambda}}(\llbracket \varphi \rrbracket^{\mathfrak{M}}))) \\ &= \theta'_{upX}(\llbracket \heartsuit^{\lambda} \varphi \rrbracket^{\mathfrak{M}}) \\ &= \llbracket \underbrace{\heartsuit^{\lambda} \varphi}^{\mathbb{M}} \underbrace{\blacksquare}^{\mathfrak{M}} \end{split}$$

This implies

$$\mathfrak{q} \Vdash \heartsuit^{\lambda} \varphi \quad \text{iff} \quad \mathfrak{q} \in \llbracket \heartsuit^{\lambda}(\varphi) \rrbracket^{pf\mathfrak{M}} \quad \text{iff} \quad \mathfrak{q} \in \llbracket \heartsuit^{\lambda} \varphi \rrbracket^{\mathfrak{M}} \quad \text{iff} \quad \llbracket \heartsuit^{\lambda} \varphi \rrbracket^{\mathfrak{M}} \in \mathfrak{q}$$

The case for n-ary modalities is analogous.

5.26 Corollary. Let \mathfrak{M} be a \mathcal{T} -model and $x \in \mathfrak{M}$. Then for all $\varphi \in \mathbb{L}(\Lambda)$ we have

$$\mathfrak{M}, x \Vdash \varphi \quad iff \quad \widehat{\mathfrak{M}}, \widetilde{x} \Vdash \varphi$$

Proof. We have $x \Vdash \varphi$ iff $x \in [\![\varphi]\!]^{\mathfrak{M}}$ iff $[\![\varphi]\!]^{\mathfrak{M}} \in \widetilde{x}$ iff $\widetilde{x} \Vdash \varphi$. The last "iff" follows from Proposition 5.25.

5.27 Corollary. Let $\mathfrak{M} = (X, \gamma, V)$ be a \mathcal{T} -model. Then for all $\varphi \in \mathbb{L}(\Lambda)$ we have

$$\mathfrak{M}\Vdash\varphi\quad iff\quad \mathfrak{\widehat{M}}\Vdash\varphi.$$

Proof. The direction from right to left follows from Corollary 5.26. For the converse, observe that $\mathfrak{M} \Vdash \varphi$ implies $\llbracket \varphi \rrbracket^{\mathfrak{M}} = X$, so by definition of a prime filter we have $\llbracket \varphi \rrbracket^{\mathfrak{M}} \in \mathfrak{q}$ for all $\mathfrak{q} \in \widehat{\mathfrak{M}}$ and hence $\widehat{\mathfrak{M}} \Vdash \varphi$.

5.28 Corollary. Let \mathfrak{X} be a \mathcal{T} -frame and suppose $pf\mathfrak{X} \Vdash \varphi$. Then $\mathfrak{X} \Vdash \varphi$.

Proof. Let \mathfrak{M} be a \mathcal{T} -model based on \mathfrak{X} . Then $\widehat{\mathfrak{M}}$ is a \mathcal{T} -model based on $\widehat{\mathfrak{X}}$, so by assumption $\widehat{\mathfrak{M}} \Vdash \varphi$. It follows from Corollary 5.26 that $\mathfrak{M} \Vdash \varphi$.

6 Definability

In this section is we prove the Goldblatt-Thomason theorem for canonical dialgebraic intuitionistic logics given by predicate liftings and axioms, interpreted in \mathcal{T} -frames and \mathcal{T} -models. Just like the original proof by Goldblatt and Thomason [14], we use of Birkhoff's variety theorem to obtain an algebraic counterpart of the Goldblatt-Thomason theorem.

In subsections 6.1 and 6.2 below we proof Goldblatt-Thomason theorems for frames and models of canonical dialgebraic intuitionistic logics. In Section 7 below we show how this general result can be used to obtain Goldblatt-Thomason theorems for the modal intuitionistic logic studied by Fischer Servi [10, 11] and Plotkin and Sterling [26].

Before commencing, let us briefly recall Birkhoff's variety theorem and a version of the theorem for ground equations. A ground equation in the setting of universal algebra is an equation that does not use any variables. For a subcategory A of a category $Alg\Sigma$ of algebras of some signature Σ , we write EA, HA, SA and PA for the closure of A under embeddings (E), homomorphic images (H), subalgebras (S) and products (P), respectively.

Jim de Groot

6.1 Birkhoff Variety Theorem. Let A be a class of algebras. Then

1. A is definable by equations if and only if A = HSPA;

2. A is definable by ground equations if and only if A = EHSPA.

The first item is well known and a proof can be found in e.g. [4]; a proof sketch of the second item is given in [22, Theorem 3.13].

6.1 The Goldblatt-Thomason theorem for frames

Finally, we put our theory to work and prove a Goldblatt-Thomason theorem for modal intuitionistic logics. We let (\mathcal{L}, ρ) be some canonical logic for \mathcal{T} given by predicate liftings and axioms.

6.2 Definition. Let $\Phi \subseteq \mathbb{L}(\Lambda)$ and $\mathfrak{X} \in \mathsf{Dialg}(i, \mathcal{T})$. Recall that $(X, \gamma) \Vdash \Phi$ if $(X, \gamma) \Vdash \varphi$ for all $\varphi \in \Phi$. Define

$$\operatorname{Fr} \Phi = \{ \mathfrak{X} \in \mathsf{Dialg}(i, \mathcal{T}) \mid \mathfrak{X} \Vdash \Phi \}$$

We call a class \mathcal{K} of frames *axiomatic* if $\mathcal{K} = \operatorname{Fr} \Phi$ for some Φ .

In order to employ Birkhoff's theorem, we make the following simple observation.

6.3 Lemma. If a logic (\mathcal{L}, ρ) is given by predicate liftings and axioms, then the category $\text{Dialg}(\mathcal{L}, j)$ is a variety of algebras.

Proof. We already know that the category HA of Heyting algebras is a variety of algebras. We add to its signature an *n*-ary operation symbol for each *n*-ary predicate lifting in Λ , and to the set of equations for HA the equations obtained from Ax.

6.4 Definition. If $\Phi \subset \mathbb{L}(\Lambda)$ and $\mathcal{A} \in \mathsf{Dialg}(\mathcal{L}, j)$ then we write $\mathcal{A} \models \Phi$ if $\mathcal{A} \models \varphi$ for all $\varphi \in \Phi$. Besides, we let

$$\operatorname{Alg} \Phi = \{ \mathcal{A} \in \mathsf{Dialg}(\mathcal{L}, j) \mid \mathcal{A} \models \Phi \}$$

be the collection of (\mathcal{L}, j) -dialgebras satisfying Φ . We say that a class $A \subseteq \text{Dialg}(\mathcal{L}, j)$ is *axiomatic* if $A = \text{Alg }\Phi$ for some collection Φ of $\mathbb{L}(\Lambda)$ -formulae.

Clearly, a class of (\mathcal{L}, j) -dialgebras is axiomatic if and only if it is a variety. Indeed, if $\mathsf{A} = \{\mathcal{A} \in \mathsf{Dialg}(\mathcal{L}, j) \mid \mathcal{A} \models \Phi\}$, then it is precisely the variety of algebras satisfying $\varphi^x \leftrightarrow \top$, where $\varphi \in \Sigma$ and φ^x is the formula we get from φ by replacing the proposition letters with variables from some set S of variables. Conversely, suppose A is a variety of algebras given by a set E of equations using variables in S. For each equation $\varphi = \psi$ in E, let $(\varphi \leftrightarrow \psi)^p$ be the formula we get from replacing the variables in $\varphi \leftrightarrow \psi$ with proposition letters. Then we have $\mathsf{A} = \mathsf{Alg}\{(\varphi \leftrightarrow \psi)^p \mid \varphi = \psi \in E\}.$

For a class \mathcal{K} of (i, \mathcal{T}) -dialgebras, write $\mathcal{K}^+ = \{\mathfrak{X}^+ \mid \mathfrak{X} \in \mathcal{K}\}$ for the collection of corresponding complex algebras. Then we have:

6.5 Lemma. A class $\mathcal{K} \subseteq \text{Dialg}(i, \mathcal{T})$ is axiomatic if and only if

$$\mathcal{K} = \{ \mathfrak{X} \in \mathsf{Dialg}(i, \mathcal{T}) \mid \mathfrak{X}^+ \in HSP(\mathcal{K}^+) \}.$$
(3)

Proof. Suppose \mathcal{K} is axiomatic, i.e., $\mathcal{K} = \operatorname{Fr} \Phi$. Then $HSP(\mathcal{K}^+) \subseteq \operatorname{Alg} \Phi$ so $\mathfrak{X}^+ \in HSP(\mathcal{K}^+)$ implies $\mathfrak{X} \in \operatorname{Fr} \Phi$. This proves \supseteq in (3). The other inclusion is obvious. Conversely, suppose (3) holds. Since $HSP(\mathcal{K}^+)$ is a variety it is of the from $\operatorname{Alg} \Phi$. It then follows that $\mathcal{K} = \operatorname{Fr} \Phi$. \Box

We now have all the ingredients to prove the Goldblatt-Thomason theorem for canonical dialgebraic intuitionistic logics.

6.6 Theorem. Let (\mathcal{L}, ρ) be a canonical logic for \mathcal{T} : Krip \rightarrow Pos given by predicate liftings and axioms. Suppose $\mathcal{K} \subseteq \mathsf{Dialg}(i, \mathcal{T})$ is closed under prime filter extensions. Then the following are equivalent:

- 1. *K* is axiomatic;
- 2. *K* is closed under disjoint unions, regular bisimilar images, and reflects prime filter extensions.
- 3. *K* is closed under disjoint unions, regular subframes, p-morphic images and reflects prime filter extensions.

Proof. The implication from 1 to 2 follows from Propositions 5.23 and 5.16 and Corollary 5.28. The implication $2 \Rightarrow 3$ follows from Proposition 5.18. So we only have to prove that 3 implies 1.

By Lemma 6.5 it suffices to prove that $\mathcal{K} = \{\mathfrak{X} \in \mathsf{Dialg}(i, \mathcal{T}) \mid \mathfrak{X}^+ \in HSP(\mathcal{K}^+)\}$. So suppose $\mathfrak{X} = (X, \gamma)$ is a \mathcal{T} -dialgebra such that $\mathfrak{X}^+ \in HSP(\mathcal{K}^+)$. Then there are $\mathfrak{Z}_i \in \mathcal{K}$ such that \mathfrak{X}^+ is the homomorphic image of a sub-dialgebra \mathcal{A} of the coproduct of \mathfrak{Z}_i^+ . In a diagram:

$$\mathfrak{X}^+ \iff \mathcal{A} \rightarrowtail \prod \mathfrak{Z}_i^+$$

Note that $\prod \mathfrak{Z}_i^+ = (\coprod \mathfrak{Y}_i)^+$. Dually, this yields

$$(\mathfrak{X}^+)_+ \xrightarrow{\text{injective}} \mathcal{A}_+ \longleftarrow \left(\left(\coprod \mathfrak{Y}_i \right)^+ \right)_+$$

We have $\coprod \mathfrak{Y}_i \in \mathcal{K}$ because \mathcal{K} is closed under coproducts, and $((\coprod \mathfrak{Y}_i)^+)_+ \in \mathcal{K}$ because \mathcal{K} is closed under prime filter extensions. Furthermore, $\mathcal{A}_+ \in \mathcal{K}$ since \mathcal{K} is closed under p-morphic images and $(\mathfrak{X}^+)_+ \in \mathcal{K}$ for \mathcal{K} is closed under generated subframes. Finally, since \mathcal{K} reflects prime filter extensions we have $\mathfrak{X} \in \mathcal{K}$.

As a consequence of Proposition 4.18 the theorem holds for all intuitionistic polynomial functors and their corresponding logics, defined in Subsection 4.4. In particular, this includes the modal intuitionistic logics studied by Wolter and Zakharyaschev [35, 36, 37] and intuitionistic logic with a geometric modality, studied by Goldblatt [12, Section 6].

6.2 Goldblatt-Thomason theorem for models

Our next goal is to prove the Goldblatt-Thomason theorem for models. The proof of this is similar to the proof of Theorem 6.6, but uses the variation of Birkhoff's theorem given in the second item of Birkhoff's variety theorem (Theorem 6.1).

Let us fix some notation. Let (\mathcal{L}, ρ) be a canonical logic for $\mathcal{T} : \mathsf{Krip} \to \mathsf{Pos}$ given by a pair $(\Lambda, \operatorname{Ax})$ of predicate liftings and axioms. For $\Phi \subseteq \mathbb{L}(\Lambda)$, let

$$\operatorname{Mod} \Phi = \{\mathfrak{M} \in \mathsf{Mod}(\mathcal{T}) \mid \mathfrak{M} \Vdash \Phi\}.$$

We say that $\mathcal{K} \subseteq \mathsf{Mod}(\mathcal{T})$ is *axiomatic* if it is for the form $\mathrm{Mod}\,\Phi$ for some set Φ of $\mathbb{L}(\Lambda)$ -formulae. Similarly, for algebraic models we set

$$\operatorname{AlgMod} \Phi = \{ \mathcal{A} \in \mathsf{AlgMod}(\mathcal{L}) \mid \mathcal{A} \models \Phi \}$$

and call $A \subseteq AlgMod(\mathcal{L})$ axiomatic if it is of the form $AlgMod \Phi$.

Like Lemma 6.3, we can make the observation that $\mathsf{AlgMod}(\mathcal{L})$ is a variety whenever \mathcal{L} is given by predicate liftings and relations. Indeed, we already know that $\mathsf{Dialg}(\mathcal{L}, j)$ is a variety of algebras, and we can simply add to this a nullary operator corresponding to each proposition letter $p \in \mathsf{Prop}$.

Let us derive an analog of Lemma 6.5. Intuitively, we now think of the proposition letters as constants in the signature of our algebra, rather than variables. As a consequence, a collection $\Phi \subseteq \mathbb{L}(\Lambda)$ corresponds to a collection of ground equations for AlgMod(\mathcal{L}).

6.7 Lemma. M is axiomatic if and only if

$$\mathcal{M} = \{\mathfrak{M} \in \mathsf{Mod}(\mathcal{T}) \mid \mathfrak{M}^+ \in EHSP(\mathcal{K}^+)\}.$$
(4)

Proof. If \mathfrak{M} is axiomatic then it is of the form $\mathfrak{M} = \operatorname{Mod} \Phi$. Consequently $EHSP(\mathfrak{M}^+) \subseteq$ AlgMod Φ so $\mathfrak{M}^+ \in EHSP(\mathfrak{M}^+)$ implies $\mathfrak{M}^+ \in \operatorname{AlgMod}(\Phi)$, which in turn implies $\mathfrak{M} \in$ Mod $\Phi = \mathfrak{M}$. The converse inclusion of (4) is obvious. Conversely, if (4) holds then \mathfrak{M}^+ is definable by a collection of ground equations Φ , and it is easy to see that $\mathfrak{M} = \operatorname{Mod} \Phi$. \Box

6.8 Theorem. Let (\mathcal{L}, ρ) be a canonical logic for \mathcal{T} : Krip \rightarrow Pos given by a pair (Λ, Ax) of predicate liftings and axioms, and let \mathcal{M} be a class of \mathcal{T} -models. Then the following are equivalent:

- 1. M is axiomatic;
- 2. M is closed under disjoint unions, m-bisimilar images, and invariant under prime filter extensions;
- 3. M is closed under disjoint unions, p-morphic domains, p-morphic images, and invariant prime filter extensions.

Proof. The implications $1 \Rightarrow 2 \Rightarrow 3$ follow from Propositions 5.22, 5.12 and 5.19 and Corollary 5.27. The proof of the implication $3 \Rightarrow 1$ is similar to the analogous implication in Theorem 6.6.

Again, the theorem holds for the intuitionistic polynomial functors and their logic from Subsection 4.4.

7 Adaptation for FS

A well-known modal intuitionistic logic, studied by e.g. Fischer Servi [10, 11] and Plotkin and Sterling [26] is the extension of WZ from Example 4.2 with the axioms

- $(FS_1) \diamond (p \to q) \to (\Box p \to \diamond q)$
- (FS₂) $(\Diamond p \to \Box q) \to \Box (p \to q)$

We will denote this extension by **FS**. We denote the language that arises from enriching intuitionistic logic with unary modal operators \Box and \diamond by $\mathbb{L}_{\Box\diamond}$, that is, $\mathbb{L}_{\Box\diamond}$ is defined by the grammar

 $\varphi ::= p \mid \top \mid \bot \mid \varphi \land \varphi \mid \varphi \lor \varphi \mid \varphi \to \varphi \mid \Box \varphi \mid \Diamond \varphi.$

In [11] Fischer Servi provides frame semantics for FS by means of *birelational frames*, and the same semantics were used in [26].

In this section we develop a Goldblatt-Thomason theorem for the logic \mathbf{FS} interpreted in birelational frames and models. First, in Subsection 7.1 we recall the definition of birelational frames and models, and the interpretation of formulae in them. We equip the classes of birelational frames and models with suitable (truth-preserving) notions of morphisms so that they become categories. We also define disjoint unions and prime filter extensions, so that we can state the Goldblatt-Thomason theorems for \mathbf{FS} in Theorems 7.10 and 7.11.

Thereafter, in Subsection 7.2 we show how birelational frames correspond to a certain class of WZ-frames. We then use this correspondence to prove the Goldblatt-Thomason theorem for frames in Subsection 7.3, leaving the proof of the analogous theorem for models to the reader.

7.1 Fischer-Servi semantics

7.1 Definition. A *BR-frame* is a tuple (X, \leq, S) where (X, \leq) is a pre-order and S is a binary relation on X that satisfies the following properties:

(BR₁) If $x \leq y$ and xSz then there exists $w \in X$ such that ySw and $z \leq w$;

(BR₂) If xSz and $z \leq w$ then there exists $y \in X$ such that $x \leq y$ and ySw.

In diagrams:

A *BR*-model is a tuple (X, \leq, S, V) consisting of an BR-frame (X, \leq, S) and a valuation V: Prop $\rightarrow \mathcal{U}p(X, \leq)$. (Here $\mathcal{U}p(X, \leq)$ denotes the collection of up-closed subsets of the pre-order (X, \leq) .) The interpretation of a formula $\varphi \in \mathbb{L}_{\Box \diamond}$ in a state x of a BR-model $\mathfrak{M} = (X, \leq, S, V)$ is defined recursively. The clauses for proposition letters and intuitionistic connectives are as in Definition 2.5, and

$$\begin{array}{ll}\mathfrak{M}, x \Vdash \Box \varphi & \text{iff} \quad x \leq ySz \text{ implies } \mathfrak{M}, z \Vdash \varphi \\ \mathfrak{M}, x \Vdash \diamond \varphi & \text{iff} \quad \exists y \in X \text{ s.t. } xSy \text{ and } \mathfrak{M}, y \Vdash \varphi \end{array}$$

Observe that this is different from the interpretation of \Box in \Box -frames because there is an additional " \leq " in the definition of its interpretation.

An immediate difference between BR-frames and the theory we have developed in the previous sections is that (X, \leq) is a pre-order rather than a poset. However, this poses no real problem as every BR-frame and model can be turned into a BR-frame or model based on a poset that satisfies precisely the same formulae. Before we make this precise we make a second observation: For a BR-frame (X, \leq, S) let the *convexification* of S be the relation \check{S} on X defined by $X\check{S}y$ iff there are $z, z' \in X$ such that xSz and xSz' and $z \leq y \leq z'$. Then we can replace S by \check{S} without affecting truth of formulae at any state.

We now define the *convexification* of a frame, which simultaneously changes underlying pre-order into a poset and makes the relation S convex.

7.2 Definition. Suppose given a BR-frame $\mathfrak{X} = (X, \leq, S)$. Let (\overline{X}, \preceq) be the poset obtained from (X, \leq) by identifying cycles. That is, we set $\overline{X} = X/\sim$ where \sim is the equivalence relation on X defined by $x \sim y$ if $x \leq y$ and $y \leq x$. Writing \overline{x} for the equivalence class of x in \overline{X} , we then set $\overline{x} \leq \overline{y}$ if $x \leq y$. It is clear that this does not depend on the choice of representative.

Define the binary relation \overline{S} on \overline{X} by $\overline{x}\overline{S}\overline{y}$ if there are $z, z' \in X$ such that xSz and xSz'and $z \leq y \leq z'$. In other words, the successor-set of \overline{x} is given by

$$\bar{S}[\bar{x}] = \mathcal{C}on_{\prec}\{\bar{y} \mid xSy\}.$$

Here $\mathcal{C}on_{\preceq}A$ denotes the smallest convex subset containing $A \subseteq \overline{X}$ with respect to the order \preceq . We call $\overline{\mathfrak{X}} = (\overline{X}, \preceq, \overline{S})$ the *convexification* of \mathfrak{X} . A BR-frame \mathfrak{X} is called *convex* if $\mathfrak{X} = \overline{\mathfrak{X}}$.

The convexification of an BR-model $\mathfrak{M} = (X, \leq, S, V)$ is defined to be $\overline{\mathfrak{M}} = (\overline{X}, \preceq, \overline{S}, \overline{V})$, where $\overline{V}(p) = \{\overline{x} \mid x \in V(p)\}$. We call an BR-model \mathfrak{M} convex if $\mathfrak{M} = \overline{\mathfrak{M}}$.

7.3 Lemma. The convexification of a BR-frame is again a BR-frame.

Proof. Let (X, \leq, S) be an BR-frame and (\bar{X}, \leq, \bar{S}) its convexification. We need to verify that it satisfies the fill-in conditions from Definition 7.1. We prove the first one, the second being similar. Suppose $\bar{x} \leq \bar{y}$ and $\bar{x}\bar{S}\bar{w}$. Then $x \leq y$ and by definition of \bar{S} there exists $v \in X$ such that xSv and $w \leq v$. Since (X, \leq, S) is an BR-frame there exists $z \in X$ such that ySz and $v \leq z$. This implies $\bar{y}\bar{S}\bar{z}$ and $\bar{w} \leq \bar{v} \leq \bar{z}$, so $\bar{w} \leq \bar{z}$, as desired.

7.4 Proposition. Let $\mathfrak{M} = (X, \leq, S, V)$ be a BR-model, $\overline{\mathfrak{M}} = (\overline{X}, \preceq, \overline{S}, \overline{V})$ its convexification, and $x \in X$. Then we have

$$\mathfrak{M}, x \Vdash \varphi \quad iff \quad \overline{\mathfrak{M}}, \overline{x} \Vdash \varphi$$

for all $x \in X$ and $\varphi \in \mathbb{L}_{\Box \diamond}$.

Proof. The proof proceeds by induction on the structure of φ . If φ is a proposition letter then the statement follows from the fact that $x \in V(p)$ if and only if $\bar{x} \in \bar{V}(p)$, which in turn follows from the fact that V(p) is up-closed in (X, \leq) . The cases for \bot, \top, \lor, \land and \rightarrow are straightforward.

Suppose $\varphi = \diamond \psi$. If $\mathfrak{M}, x \Vdash \diamond \psi$ then there exists $y \in X$ such that xSy and $\mathfrak{M}, y \Vdash \psi$. By construction of \overline{S} and induction this implies $\overline{x}\overline{S}\overline{y}$ and $\overline{\mathfrak{M}}, \overline{y} \Vdash \psi$ so that $\overline{\mathfrak{M}}, \overline{x} \Vdash \diamond \psi$. Conversely, if $\overline{\mathfrak{M}}, \overline{x} \Vdash \diamond \psi$ then there exists $\overline{y} \in \overline{X}$ such that $\overline{x}\overline{S}\overline{y}$ and $\overline{\mathfrak{M}}, \overline{y} \Vdash \psi$. By induction we have $\mathfrak{M}, y \Vdash \psi$ and as a consequence of the construction of \overline{S} we have $z \in X$ such that xSz and $y \leq z$. Persistence then implies $\mathfrak{M}, z \Vdash \psi$ so that z witnesses the fact that $\mathfrak{M}, x \Vdash \diamond \psi$.

A similar argument proves the inductive step for $\varphi = \Box \psi$.

7.5 Corollary. 1. Let \mathfrak{M} be an BR-model. Then $\mathfrak{M} \Vdash \varphi$ iff $\overline{\mathfrak{M}} \Vdash \varphi$.

2. Let \mathfrak{X} be an BR-frame. Then $\mathfrak{X} \Vdash \varphi$ iff $\overline{\mathfrak{X}} \Vdash \varphi$.

Proof. The first item follows immediately from Proposition 7.4. The second follows from the observation that for every BR-frame \mathfrak{X} we have a bijective correspondence between valuations for \mathfrak{X} and valuations for $\overline{\mathfrak{X}}$.

As a consequence of Corollary 7.5 axiomatic classes of BR-frames and BR-models are invariant under convexification. We briefly define the constructions needed to state the Goldblatt-Thomason theorem for **FS**, interpreted in BR-frames and -models.

7.6 Definition. A BR-frame morphism from (X, \leq, S) to (X', \leq', S') is a bounded morphism $f: (X, \leq) \to (X', \leq')$ such that for all $x, y \in X$ and $z' \in X'$ we have:

- 1. If xSy then f(x)S'f(y);
- 2. If f(x)S'z' then there exist $z_1, z_2 \in X$ such that $f(z_1) = f(z_2) = z'$ and $x(\leq \circ S)z_1$ and $x(S \circ \geq)z_2$.

We denote the category of BR-frames and BR-frame morphisms by BR.

A BR-model morphism $f: (X, \leq, S, V) \to (X', \leq', S', V')$ is a BR-frame morphism between the underlying BR-frames that additionally satisfies $V = f^{-1} \circ V$. The category of BR-models and BR-model morphisms is denoted by BRMod.

The definition of a BR-frame morphism may look slightly strange: it has two "backconditions". We could simply have stipulated that f(x)Sz' implies the existence of some $z \in X$ such that xSz and f(z) = z'. While this certainly preserves truth, it turns out to be slightly too restrictive for our purpose of obtaining a Goldblatt-Thomason theorem from the results from the previous sections, as it complicates the connection between convex BR-frames and certain WZ-frames (see Subsection 7.2 below).

7.7 Definition. A BR-frame $\mathfrak{B} = (X, \leq, S)$ is called a *regular subframe* of a BR-frame $\mathfrak{B}' = (X', \leq', S)$ if there exists a BR-frame morphism $m : \mathfrak{B} \to \mathfrak{B}'$ such that that $m : (X, \leq) \to (X', \leq')$ is an embedding. The BR-frame \mathfrak{B} is a *BR-morphic image* of \mathfrak{B}' if there exists a surjective BR-frame morphism $e : \mathfrak{B}' \to \mathfrak{B}$.

Let $f : \mathfrak{M} \to \mathfrak{M}'$ be a BR-model morphism. Then \mathfrak{M} is called a *BR-morphic domain* of \mathfrak{M}' . If f is surjective, \mathfrak{M}' is called a *BR-morphic image of* of \mathfrak{M} .

7.8 Definition. Let $\mathfrak{X}_i = (X_i, \leq_i, S_i)$ be a family of BR-frames. The *disjoint union* of these is defined by $\coprod \mathfrak{X}_i = \mathfrak{X} = (X, \leq, S)$ where (X, \leq) is the disjoint union of the pre-orders (X_i, \leq_i) and for $x_i \in X_i$ and $y \in X$, $x_i S y$ iff $x_i S y$. It is easy to verify that this is again an BR-frame.

The disjoint union of BR-models $\mathfrak{M}_i = (X_i, \leq_i, S_i, V_i)$ is $\coprod \mathfrak{M}_i = (X, \leq, S, V)$, where (X, \leq, S) is the disjoint union of the underlying frames and V is defined by $V(p) = \bigcup V_i(p)$.

7.9 Definition. The prime filter extension of a convex BR-frame $\mathfrak{X} = (X, \leq, S)$ is $\widehat{\mathfrak{X}} = (\widehat{X}, \subseteq, \widehat{S})$, where \widehat{X} is the collection of prime filters on (X, \leq) and \widehat{S} is defined by $\mathfrak{p}\widehat{S}\mathfrak{q}$ if for all $b \in up(X, \leq)$ we have

 $b \in \mathfrak{q}$ implies $m_{\diamond}(b) \in \mathfrak{p}$ and $m_{\Box}(b) \in \mathfrak{p}$ implies $b \in \mathfrak{q}$.

The prime filter extension of a convex BR-model $\mathfrak{M} = (X, \leq, S, V)$ is $\widehat{\mathfrak{M}} = (\widehat{X}, \subseteq, \widehat{S}, \widehat{V})$, where $\widehat{V}(p) = \{\mathfrak{q} \in \widehat{X} \mid V(p) \in \mathfrak{q}\}.$

The prime filter extension of any BR-frame or -model is defined to be the prime filter extension of its convexification.

Now that we have defined the required constructions, we can state the Goldblatt-Thomason theorems for BR-frames and BR-models.

7.10 Theorem. Let $\mathcal{K} \subseteq \mathsf{BR}$ be a class of BR-frames that is closed under prime filter extensions. Then the following are equivalent:

- 1. \mathcal{K} is axiomatic;
- 2. K is closed under disjoint unions, regular subframes and BR-morphic images, and K reflects prime filter extensions.

7.11 Theorem. Let $\mathcal{M} \subseteq \mathsf{BRMod}$ be a class of BR-models. Then the following are equivalent:

- 1. M is axiomatic;
- 2. M is closed under disjoint unions, BR-morphic domains and images, and invariant under prime filter extensions.

Jim de Groot

7.2 Connection between BR-frames and WZ-frames

In this subsection we explore the connection between BR-frames and WZ-frames. In Subsection 7.3 we use this to prove Theorem 7.10. Throughout the remainder of this section we focus on frames, rather than models. However, the proof of Theorem 7.11 is completely analogous.

Recall from Example 4.2 that a WZ-frame is a tuple $(X, \leq, R_{\Box}, R_{\diamond})$ where (X, \leq) is a poset and R_{\Box}, R_{\diamond} are binary relations on X that satisfy

$$(\leq \circ R_{\Box} \circ \leq) = R_{\Box}, \qquad (\geq \circ R_{\diamondsuit} \circ \geq) = R_{\diamondsuit}.$$

A WZ-frame morphism from $(X, \leq, R_{\Box}, R_{\diamond})$ to $(X', \leq', R'_{\Box}, R'_{\diamond})$ is a function $f : X \to X'$ that is bounded with respect to all three orders. We write WZ for the category of WZ-frames and WZ-frame morphisms. A straightforward verification shows that

$$\mathsf{WZ} \cong \mathsf{Dialg}(i, \mathcal{P}_{up} \times \mathcal{P}_{dn})$$

We equip these frames with the logic described in Example 4.2, which interprets \Box via $x \Vdash \Box \varphi$ iff $xR_{\Box}y$ implies $y \Vdash \varphi$ and $x \Vdash \Diamond \varphi$ if there exists y such that $xR_{\Diamond}y$ and $y \Vdash \varphi$.

7.12 Definition. We call a WZ-frame $(X, \leq, R_{\Box}, R_{\Diamond})$ an *FS-frame* if it satisfies:

- (P₁) If $xR_{\diamond}y$ then $\exists z \in X$ such that $y \leq z$ and $xR_{\Box}z$ and $xR_{\diamond}z$;
- (P₂) If $xR_{\Box}y$ then $\exists z \in X$ such that $x \leq z$ and $zR_{\Box}y$ and $zR_{\diamond}y$.

Diagrammatically we can depict these conditions as follows:



We write FS-WZ for the full subcategory of WZ with FS-frames as objects.

7.13 Proposition. Let $\mathfrak{X} = (X, \leq, R_{\Box}, R_{\Diamond})$ be a WZ-frame. Then we have:

- 1. $\mathfrak{X} \Vdash \Diamond (p \to q) \to (\Box p \to \Diamond q)$ if and only if \mathfrak{X} satisfies (P_1) ;
- 2. $\mathfrak{X} \Vdash (\Diamond p \to \Box p) \to \Box (p \to q)$ if and only if \mathfrak{X} satisfies (P₂).

Proof. See Subsection A.2 in the appendix.

7.14 Lemma. The class of FS-frames is closed under prime filter extensions in WZ.

Proof. Let $\mathfrak{X} = (X, \leq, R_{\Box}, R_{\diamond})$ be an FS-frame. Then the prime filter extension \mathfrak{X} is precisely the frame underlying the descriptive WZ-frame $(\hat{X}, \subseteq, \hat{R}_{\Box}, \hat{R}_{\diamond}, A)$, where $A = \{\tilde{a} \mid a \in \mathcal{U}p(X, \leq)\}$ from [37], and $\hat{R}_{\Box}, \hat{R}_{\diamond}$ are defined in Example 2.16 and Subsection 3.3 respectively. Let V be an *admissible* valuation for the descriptive frame $(\hat{X}, \subseteq, \hat{R}_{\Box}, \hat{R}_{\diamond}, A)$, i.e., a map $V : \operatorname{Prop} \to A$. Then V is of the form \widehat{W} for a valuation W for $(X, \leq, R_{\Box}, R_{\diamond})$ and as a consequence of Corollary 5.27 the model $(\widehat{X}, \subseteq, \widehat{R}_{\Box}, \widehat{R}_{\diamond}, V)$ satisfies the axioms (FS₁) and (FS₂). It then follows from [35, Proposition 5] that the underlying WZ-frame \mathfrak{X} is an FS-frame.

We now connect convex BR-frames to FS-frames. Write CBR for the full subcategory of BR whose objects are convex BR-frames.

7.15 Definition. For a convex BR-frame $\mathfrak{B} = (X, \leq, S)$ define $\mathfrak{B}^{\circ} = (X, \leq, R_{\Box}, R_{\Diamond})$, where $R_{\Box} = (\leq \circ S)$ and $R_{\Diamond} = (S \circ \geq)$. For a BR-frame morphism $f : \mathfrak{B} \to \mathfrak{B}'$ define $f^{\circ} : \mathfrak{B}^{\circ} \to (\mathfrak{B}')^{\circ}$ by $f^{\circ} = f$.

7.16 Proposition. The assignment $(\cdot)^{\circ}$ defines a functor CBR \rightarrow FS-WZ. Moreover, it is surjective on objects and full.

Proof. It follows from the coherence conditions in Definition 7.1 that \mathfrak{B}° is an FS-frame, i.e., it satisfies (\mathbf{P}_1) and (\mathbf{P}_2) .

Let $f: (X, \leq, S) \to (X', \leq', S')$ be a BR-morphism and let $R_{\Box}, R_{\diamond}, R'_{\Box}, R'_{\diamond}$ be defined as in Definition 7.15. In order to prove that f is a WZ-morphism we need to show that it is bounded with respect to all three orders, \leq, R_{\Box} and R_{\diamond} . By definition of a BR-frame we know that f is bounded with respect to \leq . We show that it is bounded with respect to R_{\Box} , boundedness for R_{\diamond} can be proven similarly. First off, we have

$$xR_{\Box}y \quad \Rightarrow \quad x(\leq \circ S)y \quad \Rightarrow \quad f(x)(\leq' \circ S')f(y) \quad \Rightarrow \quad f(x)R_{\Box}'f(y).$$

Conversely, if $f(x)R'_{\Box}z'$ then $f(x) \leq w'S'z'$ for some $w' \in X'$ and by definition of a BRmorphism we can find states $z, w, v \in X$ such that the following diagram commutes:



It follows that z is such that $x \leq vSz$, so $xR_{\Box}z$, and f(z) = z'.

To see that $(\cdot)^{\circ}$ is surjective on objects, suppose given an FS-frame $\mathfrak{X} = (X, \leq, R_{\Box}, R_{\diamond})$. Set $S = R_{\Box} \cap R_{\diamond}$, i.e., xSy iff $xR_{\Box}y$ and $xR_{\diamond}y$, and define $\mathfrak{X}_{\circ} = (X, \leq, S)$. We claim that \mathfrak{X}_{\circ} is a BR-frame such that $(\mathfrak{X}_{\circ})^{\circ} = \mathfrak{X}$.

In order to prove that (X, \leq, S) is a BR-frame we need to show that it satisfies the coherence conditions between \leq and S from Definition 7.1. Suppose $x \leq y$ and xSz. Then $yR_{\diamond}z$ so by (P₁) there exists $w \in X$ such that $z \leq w$ and ySw. Similarly one can prove the other coherence condition. By construction we have $xR_{\Box}y$ iff $x(\leq \circ S)y$ and $xR_{\diamond}y$ iff $x(S \circ \geq)y$, and therefore $(\mathfrak{X}_{\diamond})^{\circ} = \mathfrak{X}$.

Finally, to see that the assignment is full, let \mathfrak{B} and \mathfrak{B}' be two BR-frames. We already know that every BR-morphism $f: \mathfrak{B} \to \mathfrak{B}'$ is also a WZ-morphism $f: \mathfrak{B}^{\circ} \to (\mathfrak{B}')^{\circ}$. Conversely, we claim that for every WZ-morphism $g: \mathfrak{B}^{\circ} \to (\mathfrak{B}')^{\circ}$ the assignment $g_{\circ} = g: \mathfrak{B} \to \mathfrak{B}'$ is a BRmorphism. This follows immediately from the definition of a BR-morphism. Since $(g_{\circ})^{\circ} = g$ this proves that $(\cdot)^{\circ}$ is full.

7.17 Proposition. Let $\mathfrak{B} = (X, \leq, S)$ be a convex BR-frame and $\varphi \in \mathbb{L}_{\Box \Diamond}$. Then we have

$$\mathfrak{B} \Vdash \varphi \quad iff \quad \mathfrak{B}^{\circ} \Vdash \varphi.$$

Proof. Since the underlying poset structure of \mathfrak{B} and \mathfrak{B}° is the same, valuations for \mathfrak{B} correspond bijectively to valuations for \mathfrak{B}° . Let V be such a valuation. We will show that

$$(X, \leq, S, V), x \Vdash \varphi \quad \text{iff} \quad (X, \leq, R_{\Box}, R_{\diamond}, V), x \Vdash \varphi.$$
(5)

38

Jim de Groot

for all $x \in X$. Clearly this proofs the proposition.

The proof of (5) proceeds by induction on the structure of φ . The propositional case is trivial, as are the cases for all intuitionistic connectives. If $\varphi = \Box \psi$ and $x \in X$ we have

$$\begin{aligned} (X,\leq,S,V), x \Vdash \Box \psi & \text{iff} \quad x(\leq \circ S)y \text{ implies } y \Vdash \psi \\ & \text{iff} \quad xR_{\Box}y \text{ implies } y \Vdash \psi \\ & \text{iff} \quad (X,\leq,R_{\Box},R_{\diamond},V), x \Vdash \Box \psi \end{aligned}$$

In case $\varphi = \Diamond \psi$ we have

$$\begin{array}{ll} (X,\leq,S,V), x \Vdash \Diamond \psi & \text{iff} \quad \exists y \in X \text{ s.t. } xSy \text{ and } y \Vdash \psi \\ & \text{iff} \quad \exists y \in X \text{ s.t. } x(S \circ \geq)y \text{ and } y \Vdash \psi \\ & \text{iff} \quad \exists y \in X \text{ s.t. } xR_{\Diamond}y \text{ and } y \Vdash \psi \\ & \text{iff} \quad (X,\leq,R_{\Box},R_{\Diamond},V), x \Vdash \Diamond \psi \end{array}$$

The second "iff" follows from the fact that truth sets of formulae are up-closed in (X, \leq) .

7.18 Lemma. Let \mathfrak{B} and \mathfrak{B}' be two convex BR-frames. Then:

- 1. \mathfrak{B} is a regular subframe of \mathfrak{B}' iff \mathfrak{B}° is a regular subframe of $(\mathfrak{B}')^{\circ}$;
- 2. \mathfrak{B} is a BR-morphic image of \mathfrak{B}' iff \mathfrak{B}° is a WZ-morphic of $(\mathfrak{B}')^{\circ}$.

Proof. \mathfrak{B} is a regular subframe of \mathfrak{B}' if and only if there exists a BR-frame morphism $m : \mathfrak{B} \to \mathfrak{B}'$ which is an embedding on the underlying posets. Then $m^\circ : \mathfrak{B}^\circ \to (\mathfrak{B}')^\circ$ is a WZ-frame morphism that is an embedding on the underlying posets, and hence witnesses the fact that \mathfrak{B}° is a regular subframe of $(\mathfrak{B}')^\circ$. The proof of the converse is similar, as is the second item. \Box

7.19 Lemma. Let $\mathfrak{B}_i = (X_i, \leq_i, S_i)$ be a family of BR-frames. Then we have

$$\left(\coprod\mathfrak{B}_i\right)^\circ=\coprod\mathfrak{B}_i^\circ.$$

Proof. This follows from the definitions of disjoint unions in BR and WZ.

7.20 Lemma. Let $\mathfrak{B} = (X, \leq, S)$ be a convex BR-frame. Then $\widehat{\mathfrak{B}} = (\widehat{\mathfrak{B}})_{\alpha}$.

Proof. This follows from the definitions of prime filter extensions in BR and WZ.

7.3 Proof of the Goldblatt-Thomason Theorem for BR-frames

Let $\mathcal{K} \subseteq \mathsf{BR}$ be a class of BR-frames closed under prime filter extensions. We separately prove both implications of the theorem.

Proof of Theorem 7.10 $(1 \Rightarrow 2)$. If \mathcal{K} is the collection of frames validating some collection Φ of formulae, then it follows from a straightforward verification that \mathcal{K} is closed under BR-morphisms and disjoint unions. Let us show that \mathcal{K} reflects prime filter extensions. Let $\mathfrak{B} = (X, \leq, S)$ be a BR-frame with prime filter extension $\widehat{\mathfrak{B}}$ and suppose $\widehat{\mathfrak{B}} \Vdash \varphi$. It follows from Corollary 7.5 that order to prove $\mathfrak{B} \Vdash \varphi$ it suffices to prove that $\overline{\mathfrak{B}} \Vdash \varphi$. Note that $\overline{\mathfrak{B}} = \widehat{\mathfrak{B}}$ be definition, so that $\overline{\mathfrak{B}} \Vdash \varphi$. Furthermore, Lemma 7.20 shows that $\overline{\mathfrak{B}} = (\widehat{(\mathfrak{B})^\circ})_\circ$. It follows from Proposition 7.17 and Corollary 5.28 that $(\widehat{(\mathfrak{B})^\circ})_\circ \Vdash \varphi$ implies $\overline{\mathfrak{B}} \Vdash \varphi$. This ultimately proves $\mathfrak{B} \Vdash \varphi$, so \mathcal{K} reflects prime filter extensions.

Jim de Groot

Proof of Theorem 7.10 $(2 \Rightarrow 1)$. Now suppose \mathcal{K} is closed under disjoint unions, regular subframes, BR-morphic images and \mathcal{K} reflects prime filter extensions. Since then \mathcal{K} preserves and reflects prime filter extensions, the class \mathcal{K} is invariant under convexification.

Consider the class

 $\mathcal{K}^{\circ} = \{\mathfrak{B}^{\circ} \mid \mathfrak{B} \in \mathsf{CBR} \text{ and } \mathfrak{B} \in \mathcal{K}\}.$

It follows from Lemmas 7.18, 7.19 and 7.20, and the fact FS-WZ is axiomatic, that \mathcal{K}° is closed under disjoint unions, regular subframes, p-morphic images and that it reflects prime filter extensions. Therefore, as a consequence of Theorem 6.6 it is axiomatic, so there exists a collection $\Phi \subseteq \mathbb{L}_{\Box \diamond}$ of formulae such that $\mathcal{K}^{\circ} = \operatorname{Fr} \Phi$. As a consequence we have

$\mathfrak{B}\in \mathcal{K}$	iff	$ar{\mathfrak{B}}\in \mathscr{K}$	$(\mathcal R \text{ invariant under convexification})$
	iff	$\bar{\mathfrak{B}}^\circ\in \mathscr{K}^\circ$	(Definition of \mathcal{K}°)
	iff	$\bar{\mathfrak{B}}^{\circ} \Vdash \Phi$	$(\mathscr{K}^\circ = \operatorname{Fr} \Phi)$
	iff	$\bar{\mathfrak{B}} \Vdash \Phi$	(Proposition 7.17)
	iff	$\mathfrak{B}\Vdash\varphi$	(Corollary 7.5)

So $\mathcal{K} = \{\mathfrak{B} \in \mathsf{BR} \mid \mathfrak{B} \Vdash \Phi\}.$

The proof of Theorem 7.11 is similar, and is left to the reader.

8 Conclusion

We have given Goldblatt-Thomason theorems for frames and models of a large class of modal intuitionistic logics, including ones studied by Wolter and Zakharyaschev [35, 36, 37], Goldblatt [12, Section 6], Fischer Servi [11], and Plotkin and Sterling [26]. There are many avenues for further research, some of which are listed below.

- More canonical logics An immediate question is whether there are more examples of canonical dialgebraic intuitionistic logics. One such example can be found in intuitionistic logic with a constructive strict implication [24], although no details have been provided in this paper. With each additional canonical dialgebraic intuitionistic logic we can enlarge the class of intuitionistic polynomial functors and corresponding logics for which the Goldblatt-Thomason theorems hold.
- More Goldblatt-Thomason theorems It is reasonable to expect that similar techniques as the ones used in Section 7 give rise to Goldblatt Thomason theorems for other modal intuitionistic logics, like the ones studied by Božić and Došen [3].
- **Bi- and dual-intuitionistic logic** The results of this paper can be adapted to the setting of modal bi-intuitionistic logic (see e.g. [15]) and modal dual-intuitionistic logic in a straightforward manner.
- Adaptation to other base logics Elaborating on the previous item, one may wonder to what extend the techniques in this paper generalise to the general setting of dialgebraic modal logic presented in [17]. This road for further research is particularly attractive because dialgebraic logic encompasses coalgebraic logic, so a general result for dialgebras would carry over to the realm of coalgebraic modal logic.

Descriptive frames Each of the logics described in the examples of Section 2 and Section 3 can be given a notion of descriptive frame, which is closely related to the definition of τ . In each case we can describe these descriptive frames as a category of dialgebras for the inclusion $\mathcal{I} : \mathsf{ES} \to \mathsf{Pries}$ of Esakia spaces into Priestley spaces. Adapting techniques from Section 4 one can probably obtain such a description of descriptive frames for all logics corresponding to the intuitionistic polynomial functors defined in Subsection 4.4 in a similar fashion as [20, Section 4].

References

- J. Barwise and L. S. Moss. Modal correspondence for models. Journal of Philosophical Logic, 27:275–294, 1998.
- [2] P. Blackburn, M. de Rijke, and Y. Venema. *Modal Logic*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, Cambridge, 2001.
- [3] M. Božić and K. Došen. Models for normal intuitionistic modal logics. Studia Logica, 43:217–245, 1984.
- [4] S. Burris and H. P. Sankappanavar. A Course in Universal Algebra. Springer, 1981.
- [5] S. A. Celani and R. Jansana. Priestley duality, a Sahlqvist theorem and a Goldblatt-Thomason theorem for positive modal logic. *Logic Journal of the IGPL*, 7:683–715, 1999.
- [6] A. Chagrov and M. Zakharyaschev. Modal Logic. Oxford University Press, Oxford, 1997.
- [7] B. F. Chellas. Modal Logic: An Introduction. Cambridge University Press, Cambridge, 1980.
- [8] W. Conradie, A. Palmigiano, and A. Tzimoulis. Goldblatt-thomason for LE-logics, 2018.
- [9] L. L. Esakia. *Heyting Algebras*. Trends in Logic. Springer, Springer, 2019. Translated by A. Evseev.
- [10] G. Fischer Servi. On modal logic with an intuitionist base. Studia Logica, pages 141–149, 1977.
- [11] G. Fischer Servi. Semantics for a class of intuitionistic modal calculi. In D. Chiara and M. Luisa, editors, *Italian Studies in the Philosophy of Science*, pages 59–72, Dordrecht, 1981. Springer Netherlands.
- [12] R. I. Goldblatt. Mathematics of Modality. CSLI publications, Stanford, California, 1993.
- [13] R. I. Goldblatt. Axiomatic classes of intuitionistic models. Journal of Universal Computer Science, 11(12):1945–1962, 2005.
- [14] R. I. Goldblatt and S. K. Thomason. Axiomatic classes in propositional modal logic. In J. Crossley, editor, Algebra and Logic, pages 163–173. Springer, 1974.
- [15] J. de Groot and D. Pattinson. Hennessy-Milner properties for (modal) bi-intuitionistic logic. In R. Iemhoff, M. Moortgat, and R. de Queiroz, editors, *Proc. WoLLIC 2019*, pages 161–176, Berlin, Heidelberg, 2019. Springer.
- [16] J. de Groot and D. Pattinson. Hennessy-Milner properties via compactness, 2020. Under review, submitted March 2020.
- [17] J. de Groot and D. Pattinson. Modal intuitionistic logics as dialgebraic logics. In Proc. LICS'20, pages 355—369, New York, 2020. Association for Computing Machinery.
- [18] H. H. Hansen. Monotonic modal logics. Master's thesis, Institute for Logic, Language and Computation, University of Amsterdam, 2003.
- [19] K. Kojima. Which classical correspondence is valid in intuitionistic modal logic? Logic Journal of the IGPL, 20(1):331–348, 12 2011.
- [20] C. Kupke, A. Kurz, and Y. Venema. Stone coalgebras. Theoretical Computer Science, 327(1):109– 134, 2004. Selected Papers of CMCS '03.
- [21] C. Kupke and D. Pattinson. Coalgebraic semantics of modal logics: An overview. Theoretical Computer Science, 412(38):5070–5094, 2011. CMCS Tenth Anniversary Meeting.

- [22] A. Kurz and J. Rosický. The Goldblatt-Thomason theorem for coalgebras. In T. Mossakowski, U. Montanari, and M. Haveraaen, editors, *Algebra and Coalgebra in Computer Science*, pages 342–355, Berlin, Heidelberg, 2007. Springer Berlin Heidelberg.
- [23] T. Litak. Constructive modalities with provability smack, 2017.
- [24] T. Litak and A. Visser. Lewis meets Brouwer: Constructive strict implication. Indagationes Mathematicae, 29(1):36–90, 2018.
- [25] D. Pattinson. Coalgebraic modal logic: soundness, completeness and decidability of local consequence. *Theoretical Computer Science*, 309(1):177–193, 2003.
- [26] G. Plotkin and C. Stirling. A framework for intuitionistic modal logics: Extended abstract. In Proceedings of the 1986 Conference on Theoretical Aspects of Reasoning about Knowledge, TARK '86, pages 399–406, San Francisco, CA, USA, 1986. Morgan Kaufmann Publishers Inc.
- [27] H.A. Priestley. Representation of distributive lattices by means of ordered Stone spaces. Bulletin of the London Mathematical Society, 2(2):186–190, 1970.
- [28] P. H. Rodenburg. Intuitionistic Correspondence Theory. PhD thesis, University of Amsterdam, 1986.
- [29] M. Rößiger. Coalgebras and modal logic. Electronic Notes in Theoretical Computer Science, 33:294–315, 2000.
- [30] K. Sano and M. Ma. Goldblatt-Thomason-style theorems for graded modal language. In Proc. AiML 2006, pages 330–349. College Publications, 2010.
- [31] K. Sano and J. Virtema. Characterising modal definability of team-based logics via the universal modality. Annals of Pure and Applied Logic, 170:1100–1127, 2019.
- [32] B. Teheux. Modal definability based on Łukasiewicz validity relations. Studia Logica, 104:343–363, 2016.
- [33] J. van Benthem. Modal frame classes revisited. Fundamenta Informaticae, 18:307–317, 1993.
- [34] Y. Venema. Model definability, purely modal. In J. Gerbrandy, M. Marx, M. de Rijke, and Y. Venema, editors, JFAK. Essays Dedicated to Johan van Benthem on the Occasion of his 50th Birthday. Vossiuspers, Amsterdam University Press, 1999. www.illc.uva.nl/j50.
- [35] F. Wolter and M. Zakharyaschev. On the relation between intuitionistic and classical modal logics. Algebra and Logic, 36:121–155, 1997.
- [36] F. Wolter and M. Zakharyaschev. Intuitionistic modal logics as fragments of classical bimodal logics. In E. Orlowska, editor, *Logic at Work, Essays in honour of Helena Rasiowa*, pages 168– 186. Springer–Verlag, 1998.
- [37] F. Wolter and M. Zakharyaschev. Intuitionistic modal logic. In A. Cantini, E. Casari, and P. Minari, editors, Logic and Foundations of Mathematics: Selected Contributed Papers of the Tenth International Congress of Logic, Methodology and Philosophy of Science, pages 227–238, Dordrecht, 1999. Springer Netherlands.
- [38] S. Yokota. General characterization results on intuitionistic modal propositional logics. Commentarii Mathematici Universitatis Sancti Pauli, 34(2):177–199, 1985.

A Appendix

A.1 The categories Pos and Krip

In this subsection we prove that Krip has all small limits and colimits, and that they are computed as in Pos. Recall that we write Pos for the category of posets and monotone morphisms, Krip for the subcategory of posets and *bounded* morphisms, and $i : \text{Krip} \to \text{Pos}$ for the inclusion functor. We will generally write X for a poset and denote its order by \leq_X when we need it.

A.1 Lemma. The regular monomorphisms in Pos are the embeddings.

Proof. If m is a regular mono on Pos then it follows from the construction of equalisers in Pos that m is an embedding. Conversely, suppose $m: X \to Y$ is an embedding. Let Y_1, Y_2 be two copies of Y with isomorphisms $i_1: Y \to Y_1$ and $i_2: Y \to Y_2$. and let $Z = (Y_1 + Y_2)/\sim$, where $y_1 \sim y_2$ if there exists $y \in Y$ such that $y_1 = i_1(y)$ and $y_2 = i_2(y)$. Now let $j_1: Y_1 \to Z$ and $j_2: Y_2 \to Z$ be the obvious inclusion maps. Then X arises as the equaliser of the parallel pair of morphisms $j_1i_1, j_2i_2: Y \to Z$.

A.2 Proposition. The inclusion functor $i: Krip \rightarrow Pos$ preserves and reflects all small limits.

Proof. It suffices to show that the final object, equalisers and products in Krip are computed as in Pos. All of them are computed as in Set. \Box

Towards describing the colimits in Pos and Krip, we investigate their coequalisers. We first compute the the coequaliser in Pos via PreOrd, the category of preorders and monotone maps. This is convenient because all small limits and colimits in PreOrd are computed as in Set. The adjunction between PreOrd and Pos then yields the notion of a coequaliers in Pos by modding gout loops.

In particular, the coequaliser in PreOrd of a parallel pair of morphisms $f, g : X \to Y$ between posets is the set $Q' = Y/\sim$, where \sim is the smallest equivalence relation generated by $f(x) \sim g(x)$ for all $x \in X$. Order Q' by setting $z \leq_{Q'} z'$ if there are $y, y' \in Y$ such that $y \leq_Y y'$ and f(y) = z and f(y') = z'. The coequaliser $q : Y \to Q$ of f and g in Pos is then obtained by identifying loops in Q', that is $Q = Q'/\sim_{\ell}$ where $z \sim_{\ell} z'$ iff $z \leq_Q z'$ and $z' \leq_Q z$.

Now suppose f and g are bounded morphisms. We claim that this implies that q is bounded. Clearly it suffices to prove that q' is bounded. So let $y \in Y$, $z' \in Q'$ and suppose $q'(y) \leq_{Q'} z'$. Then by definition of $\leq_{Q'}$ there are $y_1, y_2 \in Y$ such that $y_1 \leq y_2$ and $q'(y_1) = q'(y)$ and $q'(y_2) = z'$. In particular, this means that $y \sim y_1$, so there is finite collection $x_1, \ldots, x_n \in X$ such that $h_1(x_1) = y_1$ and $h_i^\circ(x_i) = h_{i+1}(x_{i+1})$, where $h_i \in \{f, g\}$ and $h_i^\circ \in \{f, g\}$ is such that $h_i \neq h_i^\circ$. Using the fact that f and g are bounded, we obtain $\hat{y}_2 \in Y$ such that $y \leq \hat{y}_2$ and $y_2 \sim \hat{y}_2$. The following diagram depicts this argument:



In particular, this means \hat{y}_2 is such that $y \leq \hat{y}_2$ and $q'(\hat{y}_2) = z'$, and hence q' is bounded. An easy verification now proves that q is also the coequaliser of f and g in Krip.

A.3 Proposition. The inclusion functor $i : Krip \rightarrow Pos$ preserves and reflects all small colimits.

Proof. It is obvious that colimits and the initial object in Krip are computed as in Krip. Furthermore, we have just seen above that coequalisers in Krip are computed as in Pos. \Box

A.4 Corollary. A bounded morphism $f: X \to Y$ in Krip is a regular mono if and only if it is an embedding.

Jim de Groot

A.2 Proof of Proposition 7.13

The proof of Proposition 7.13 is split into two lemmas.

A.5 Lemma. A WZ-frame $\mathfrak{X} = (X, \leq, R_{\Box}, R_{\Diamond})$ satisfies $\Diamond(p \to q) \to (\Box p \to \Diamond q)$ if and only if \mathfrak{X} satisfies (P_1) , i.e., whenever $xR_{\Diamond}y$ there exists $z \in X$ such that $y \leq z$ and $xR_{\Box}z$ and $xR_{\Diamond}z$.

Proof. First assume that \mathfrak{X} satisfies (P₁). Let V be any valuation for \mathfrak{X} and let $x \in X$. Suppose that $x \leq y$ and $y \Vdash \Diamond (p \to q)$. We aim to prove that $y \Vdash \Box p \to \Diamond q$, as this proves that $x \Vdash \Diamond (p \to q) \to (\Box p \to \Diamond q)$. Since $y \Vdash \Diamond (p \to q)$ there exists $w \in X$ such that $yR_{\Diamond}w$ and $w \Vdash p \to q$. Now suppose $y \leq z$ and $z \Vdash \Box p$. Then $zR_{\Diamond}w$ and by (P₁) there exists $v \in X$ such that $w \leq v$ and $zR_{\Box}v$ and $zR_{\Diamond}v$. Since $z \Vdash \Box p$ this implies $v \Vdash p$, and since $w \leq v$ and $w \Vdash p \to q$ we find $v \Vdash q$. But then we have $z \Vdash \Diamond q$, because $zR_{\Diamond}v$. Diagramatically:



This proves that $y \Vdash \Box p \to \Diamond q$ and hence $x \Vdash \Diamond (p \to q) \to (\Box p \to \Diamond q)$. Since x and V were chosen arbitrarily we find that $\mathfrak{X} \Vdash \Diamond (p \to q) \to (\Box p \to \Diamond q)$.

Now suppose \mathfrak{X} does not satisfy (P₁). There there exists $x, y \in X$ such that $xR_{\diamond}y$ and for all $z \geq y$ either $\neg(xR_{\Box}z)$ or $\neg(xR_{\diamond}z)$. Define a valuation V for \mathfrak{X} by $V(p) = R_{\Box}[x]$ and

$$V(q) = R_{\Box}[x] \cap \uparrow_{<} y = \{ s \in X \mid y \le s \text{ and } xR_{\Box}s \}.$$

Then V(p) is up-closed in (X, \leq) because $(\leq \circ R_{\Box} \circ \leq) = R_{\Box}$ and V(q) is up-closed because it is the intersection of two up-closed sets.

By construction of V we have $y \Vdash p \to q$ and hence $x \Vdash \Diamond (p \to q)$. Furthermore, by definition of V(p) we have $x \Vdash \Box p$. Since $xR_{\Box}w$ and $y \leq w$ implies $\neg (xR_{\Diamond}w)$ and $V(q) \subseteq \uparrow_{\leq} y$ we have $x \not\Vdash \Diamond q$. Therefore $x \not\Vdash \Box p \to \Diamond q$, which ultimately proves $x \not\Vdash \Diamond (p \to q) \to (\Box p \to \Diamond q)$.

A.6 Lemma. A WZ-frame $\mathfrak{X} = (X, \leq, R_{\Box}, R_{\Diamond})$ satisfies $(\Diamond p \to \Box q) \to \Box (p \to q)$ if and only if \mathfrak{X} satisfies (P_2) , i.e., whenever $xR_{\Box}y$ there exists $z \in X$ such that $x \leq z$ and $zR_{\Box}y$ and $zR_{\Diamond}y$.

Proof. Suppose \mathfrak{X} satisfies (\mathbf{P}_2) and $x \in X$. Let $x \leq y$ such that $y \Vdash \Diamond p \to \Box q$. We aim to prove that $y \Vdash \Box(p \to q)$, i.e., for all w such that $yR_{\Box}w$ we want $w \Vdash p \to q$. Let v be such that $w \leq v$ and $v \Vdash p$. Then by definition of R_{\Box} we have $yR_{\Box}v$ and as a consequence of the assumed fill-in property there exists a $z \geq y$ such that $zR_{\Box}v$ and $zR_{\diamond}v$. Therefore $z \Vdash \Diamond p$ and since $y \Vdash \Diamond p \to \Box q$ this implies $z \Vdash \Box q$. But we have $zR_{\Box}v$ and hence $v \Vdash q$. This proves that $w \Vdash p \to q$ and since w was chosen arbitrarily $y \Vdash \Box(p \to q)$. Therefore, we have ultimately proved that x satisfies the desired axiom.

Now for a converse suppose that $(X, \leq, R_{\Box}, R_{\diamond})$ does not satisfy the condition from the diagram. Then there exist states y and w such that $yR_{\Box}w$ but there is no $z \in Z$ such that $y \leq z$ and $zR_{\Box}w$ and $zR_{\diamond}w$. Pick $V(p) = \uparrow_{<}w$ and

$$V(q) = \bigcup \{ R_{\Box}[z'] \mid y \le z' \text{ and } R_{\diamond}[z'] \cap \uparrow w \neq \emptyset \}.$$

Clearly p is up-closed under \leq . The same goes for q because $R_{\Box}[z']$ is closed for any state z', as a consequence of the identity $(\leq \circ R_{\Box} \circ \leq) = R_{\Box}$. Define the interpretation of any other proposition letter to be the empty set. Then $(X, \leq, R_{\Box}, R_{\diamond}, V)$ is a \Box \diamond -model.

We claim that $y \Vdash \Diamond p \to \Box q$ while $y \not\Vdash \Box(p \to q)$. Indeed, if $y \leq z$ and $z \Vdash \Diamond p$ then $R_{\Diamond}[z] \cap \uparrow w \neq \emptyset$ because $V(p) = \uparrow w$. By definition we then have $R_{\Box}[z] \subseteq V(q)$ so that $z \Vdash \Box q$. Thus y does indeed satisfy $\Diamond p \to \Box q$. To see that $y \not\Vdash \Box(p \to q)$ it suffices to prove that $w \not\models q$. Indeed, since $w \Vdash p$ this entails that $w \not\models p \to q$ and since $yR_{\Box}w$ this then proves that $y \not\models \Box(p \to q)$. Suppose towards a contradiction that $w \Vdash q$. Then $w \in V(q)$ so there exists $z' \geq y$ such that $z'R_{\Box}w$ and $z'R_{\diamond}v$ for some $v \geq w$. But then $z'R_{\diamond}w$, because $z' \geq z'R_{\diamond}v \geq w$, a contradiction with the assumption that there exists no $z' \geq y$ such that $z'R_{\Box}w$ and $z'R_{\diamond}w$. So $w \not\models q$. Therefore $y \not\models \Box(p \to q)$ and as a consequence of reflexivity of \leq we find $y \not\models (\Diamond p \to \Box q) \to \Box(p \to q)$.